

MPP-2004-169
 December 16, 2004
 gr-qc/0412067

Classification of Static, Spherically Symmetric Solutions of the Einstein-Yang-Mills Theory with Positive Cosmological Constant

Peter Breitenlohner^a, Peter Forgács^b, Dieter Maison^a

^a *Max-Planck-Institut für Physik
 — Werner Heisenberg Institut —
 Föhringer Ring 6
 D-80805 Munich, Germany*

^b *Laboratoire de Mathématiques et Physique Théorique
 CNRS UMR6083
 Université de Tours, Parc de Grandmont
 F-37200 Tours, France*

Abstract

We give a complete classification of all static, spherically symmetric solutions of the $SU(2)$ Einstein-Yang-Mills theory with a positive cosmological constant. Our classification proceeds in two steps. We first extend solutions of the radial field equations to their maximal interval of existence. In a second step we determine the Carter-Penrose diagrams of all 4-dimensional space-times constructible from such radial pieces. Based on numerical studies we sketch a complete phase space picture of all solutions with a regular origin.

1 Introduction

In contrast to the coupling of Abelian gauge fields to gravity, the interaction of non-Abelian gauge fields with gravity yields a rich structure of interesting solutions even in the spherically symmetric sector. While the Reissner-Nordström black holes are the only static, spherically symmetric, asymptotically flat solutions of the Einstein-Maxwell equations, the non-Abelian $SU(2)$ Einstein-Yang-Mills (EYM) theory has a much larger class of such solutions. There is an infinite sequence (indexed by the number n of zeros of the Yang-Mills potential) of regular ‘soliton-type’ solutions, known as the Bartnik-McKinnon (BK) solutions [1, 2, 3]. Besides there are corresponding families of black holes [4, 5, 6, 7, 3] and ‘limiting’ solutions, when n tends to infinity [8].

If a cosmological constant, Λ , is added to the theory, the asymptotic behaviour of the regular solutions changes from Minkowskian to deSitter for $\Lambda > 0$ or Anti-deSitter for $\Lambda < 0$. Furthermore for $\Lambda > 0$ the regular solutions have a cosmological horizon.

Numerical investigations of the static, spherically symmetric EYM equations with a cosmological constant have led to a rather clear picture of the most important properties of the solutions [9, 10, 11, 12]. Let us briefly recall here some of the most relevant numerical results of Refs. [9, 10, 11]. In Ref. [9] families of solutions with a regular origin and a horizon with $n = 1, 2, \dots$ zeros were found. For $0 < \Lambda < \Lambda_c(n)$ the solutions are asymptotically deSitter. At $\Lambda = \Lambda_c(n)$ the global structure of the spatial sections of the solutions changes. The area of the 2-spheres has a maximum (‘equator’) behind the horizon and then decreases back to zero, where the solutions have a singularity, implying that the spatial sections have the topology of 3-spheres. As Λ increases the horizon and the equator exchange their positions and as Λ approaches some maximal value $\Lambda_r(n)$ the radius r_h of the horizon shrinks to zero. The solutions for $\Lambda = \Lambda_r(n)$ are globally regular with compact spatial sections of 3-sphere topology. The numerical results of Ref. [10, 11] for $\Lambda < 0$ indicate the existence of generic families of asymptotically Anti-deSitter soliton-type and black hole solutions.

Besides the numerical results there are also a few analytical ones. Unfortunately the well developed general theory of dynamical systems, although very useful for local questions does not provide the tools required to answer global questions in our case. Thus special properties of the equations have to be exploited. For negative cosmological constant Winstanley [10] has given an existence proof for asymptotically Anti-deSitter black holes with *sufficiently large* radius r_h . On the other hand for positive cosmological constant Linden [13] has proven the existence of smooth solutions with a regular origin for *sufficiently small* values of Λ . Clearly it would be desirable to extend Linden’s existence proof to the numerically found families up to the 3-sphere solutions and to black holes, as well as Winstanley’s proof to all asymptotically Anti-deSitter solutions.

Experience with the existence proofs for $\Lambda = 0$ [2, 3] suggests that it is quite important to classify the possible *global* behaviour of solutions with a regular origin or a black hole horizon. Such a classification is one of the fundamental

problems in dynamical systems, and no general method is known to solve it.

The main result of this paper is a complete classification of *all* static, spherically symmetric solutions of the EYM equations with a positive cosmological constant. One should also mention here that a partial result in this direction has been already obtained by Linden [14]. Since the field equations correspond to a rather complicated dynamical system, it is highly remarkable that a complete global classification of the solutions is possible at all. It should, however, be remarked that the numerical results show that the structure of the phase space as characterized by the dependence of the solutions on the initial conditions is rather complicated. This is in particular true inside black hole or outside cosmological horizons¹ as shown in [15, 16].

Our classification proceeds in two steps. We first extend solutions of the radial field equations to their maximal interval of existence. In a second step we determine the Carter-Penrose diagrams of all 4-dimensional space-times constructible from such radial pieces.

It may be appropriate to stress the importance of an interplay between analytical and numerical methods. On the one hand it seems rather hopeless to extract sufficient information on the complex structure of the phase space just from studying the field equations analytically. On the other hand precise and reliable numerics in particular concerning ‘critical’ solutions is impossible without some analytical understanding of the behaviour of the solutions near singular points.

In Section 2 we set up the field equations. It turns out that through the introduction of an auxiliary variable they take the form of a system of coupled Riccati equations. In Section 3 we study all singular points and relate them to fixed points (f.p.s) of the dynamical system. This requires the introduction of different independent and dependent variables in each case, allowing to prove the existence and uniqueness of local solutions. In Section 4 we show that any solution starting at an arbitrary regular point and integrated in both directions either ends at one of the singular points described in Section 3 or runs towards $r = 0$ performing a kind of quasi-periodic motion around a repulsive focal point as described in [15, 16]. In Section 5 we show how to glue together 2d space-times corresponding to solutions of Eqs. (8) interpolating between two singular points and to construct their Carter-Penrose diagrams describing 4d inextendible (geodesically complete or singular) space-times.

Finally, Section 6 contains numerical results illustrating various types of solutions. In particular we illustrate how all cases of our classification fit into a complete phase space picture.

The Appendix contains a proposition stating local existence and uniqueness of solutions of a class of dynamical systems, allowing to parametrize the solutions in terms of values at f.p. In addition we formulate several Lemmata used in the proof of the classification theorems.

¹i.e. in the region where the Killing time is a space-like variable

2 Preliminaries

The line element of a spherically symmetric space-time can be written as

$$ds^2 = ds_2^2 - r^2 d\Omega^2 , \quad (1)$$

where ds_2^2 is the metric on the 2d orbit space factorizing out the action of the rotation group and $d\Omega^2$ the invariant line element of S^2 . The 2d metric can always be brought to the diagonal form

$$ds_2^2 = \sigma \left(e^{2\nu} dt^2 - e^{2\lambda} dR^2 \right) , \quad (2)$$

where $\sigma = \pm 1$ refers to regions, where t is a time-like resp. space-like coordinate. Since we consider time-independent solutions it is natural to choose t to be the Killing time. This means that we can perform a further dimensional reduction to a 1d theory. The metric functions ν , λ and r then only depend on the radial coordinate R . The latter can be chosen arbitrarily exhibiting λ as a gauge degree of freedom.

For the $SU(2)$ Yang-Mills field W_μ^a we use the standard minimal spherically symmetric (purely ‘magnetic’) ansatz

$$W_\mu^a T_a dx^\mu = W(R)(T_1 d\theta + T_2 \sin \theta d\varphi) + T_3 \cos \theta d\varphi , \quad (3)$$

where T_a denote the generators of $SU(2)$. The reduced EYM action including a cosmological constant Λ can be expressed (in suitable units) as

$$S = - \int dR e^{(\nu+\lambda)} \left[\frac{1}{2} \left(\sigma + e^{-2\lambda} ((r')^2 + \nu' (r^2)') \right) - e^{-2\lambda} W'^2 - \sigma \left(\frac{(W^2 - 1)^2}{2r^2} + \frac{\Lambda r^2}{2} \right) \right] . \quad (4)$$

In order to write the Euler-Lagrange equations in first order form we introduce

$$N \equiv e^{-\lambda} r' , \quad \kappa \equiv r e^{-\lambda} \nu' + N , \quad U \equiv e^{-\lambda} W' , \quad (5)$$

and obtain the equations of motion

$$r e^{-\lambda} N' = (\kappa - N) N - 2U^2 , \quad (6a)$$

$$r e^{-\lambda} \kappa' = \sigma (1 - 2\Lambda r^2) + 2U^2 - \kappa^2 , \quad (6b)$$

$$r e^{-\lambda} U' = \sigma W T + (N - \kappa) U , \quad (6c)$$

where

$$T = \frac{W^2 - 1}{r} . \quad (7)$$

In addition we have the only remaining diffeomorphism constraint Eq. (9) below.

We still have to fix the gauge, i.e. choose a radial coordinate. A simple choice is to use the geometrical radius r for R (Schwarzschild coordinates), yielding $e^{-\lambda} = N$. Putting $\mu \equiv \sigma N^2$, $A \equiv e^\nu / N$, and $\Lambda = 0$ Eqs. (6) become equal to Eqs. (6) in [3]. This coordinate choice has the disadvantage that the

equations become singular at stationary points of r . This problem is avoided using the gauge $e^\lambda = r$ introduced in [3], which will also be used here. In order to stress the dynamical systems character of Eqs. (5,6) we denote the corresponding radial coordinate by τ and τ derivatives by a dot. We thus get

$$\dot{r} = rN, \quad (8a)$$

$$\dot{W} = rU, \quad (8b)$$

$$\dot{N} = (\kappa - N)N - 2U^2, \quad (8c)$$

$$\dot{\kappa} = \sigma(1 - 2\Lambda r^2) + 2U^2 - \kappa^2, \quad (8d)$$

$$\dot{U} = \sigma WT + (N - \kappa)U, \quad (8e)$$

and the constraint (invariant hypersurface)

$$2\kappa N - N^2 = 2U^2 + \sigma(1 - T^2 - \Lambda r^2). \quad (9)$$

The metric function ν obeys the equation

$$\dot{\nu} = \kappa - N, \quad (10)$$

that can be solved once the other functions are known. Using the constraint Eq. (9), the equation for N can also be written as

$$\dot{N} = \frac{\sigma}{2}(1 - \Lambda r^2 - T^2) - \frac{1}{2}N^2 - U^2. \quad (11)$$

For $\Lambda = 0$ and $\sigma = +1$ these equations coincide with the Eqs. (49,50) of [3].

The quantity T , introduced in Eq. (7) as an abbreviation, obeys

$$\dot{T} = 2UW - NT. \quad (12)$$

Viewing T as an additional dependent variable we obtain a system of coupled Riccati equations (8) and (12) with two algebraic constraints. This could explain, why solutions of this system stay bounded except for simple poles at finite τ -values (as is true for a single Riccati equation).

An important quantity is the ‘mass function’

$$m = \frac{r}{2} \left(1 - \frac{\Lambda r^2}{3} - \sigma N^2 \right), \quad (13a)$$

obeying the equation

$$\dot{m} = \left(\sigma U^2 + \frac{T^2}{2} \right) rN. \quad (13b)$$

Furthermore there is a kind of τ -dependent ‘energy’ for the dynamical system Eqs. (8)

$$E = \frac{r^2}{4} \left(2\kappa N - N^2 - \sigma(1 - \Lambda r^2) \right) \quad (14a)$$

$$= \frac{\dot{W}^2}{2} - \sigma \frac{(W^2 - 1)^2}{4}, \quad (14b)$$

obeying

$$\dot{E} = (2N - \kappa)\dot{W}^2, \quad (14c)$$

which will prove useful as a ‘Lyapunov Function’.

Due to the presence of the cosmological constant the vacuum solution obtained for $W^2 = 1$ is no longer Minkowski space but deSitter space

$$ds^2 = \mu dt^2 - \frac{dr^2}{\mu} - r^2 d\Omega^2, \quad (15)$$

with

$$\mu_{\text{ds}}(r) = 1 - \frac{\Lambda r^2}{3}. \quad (16)$$

Because the radial direction changes from space-like to time-like at the cosmological horizon $r_c = \sqrt{3/\Lambda}$ we have to subdivide the radial domain into the intervals $0 < r < r_c$ and $r_c < r < \infty$ if we use the variables of Eqs. (8). The required sign change of σ going from one domain to the other can be achieved by the substitution $(d\tau, N, \kappa, U) \rightarrow (-id\tau, iN, i\kappa, iU)$.

The counterpart of the Schwarzschild solution for $\Lambda > 0$ is the Schwarzschild-deSitter (SdS) solution Eq. (15) with

$$\mu_{\text{SdS}}(r) = 1 - \frac{2M}{r} - \frac{\Lambda r^2}{3}, \quad (17)$$

describing a black hole inside a cosmological horizon² as long as μ_{SdS} has two positive zeros, i.e. $M < 1/3\sqrt{\Lambda}$. For $M = 1/3\sqrt{\Lambda}$ the function μ_{SdS} has a double zero at $r = 1/\sqrt{\Lambda}$. Geometrically this is the position of an extremal (degenerate) cosmological horizon. It corresponds to the fixed point of Eqs. (8) $(W, U, r, N, \kappa) = (\pm 1, 0, 1/\sqrt{\Lambda}, 0, \pm 1)$. As for the extremal Reissner-Nordström solution the horizon is at infinite geodesic distance in the space-like direction.

Likewise there is the Reissner-Nordström-deSitter (RNdS) solution for $W = 0$ with

$$\mu_{\text{RN}}(r) = 1 - \frac{2M}{r} + \frac{1}{r^2} - \frac{\Lambda r^2}{3}, \quad (18)$$

carrying an abelian magnetic charge. For $\Lambda > 1/4$ the function μ_{RN} has only one positive zero, while for $\Lambda < 1/4$ it has three positive zeros as long as $M_-(\Lambda) < M < M_+(\Lambda)$ for certain functions $M_{\pm}(\Lambda)$. For $M = M_-(\Lambda)$ the two smaller zeros merge to a minimum of μ_{RN} at $r_-^2 = (1 - \sqrt{1 - 4\Lambda})/2\Lambda$, while for $M = M_+(\Lambda)$ the two larger ones merge to a maximum of μ_{RN} at $r_+^2 = (1 + \sqrt{1 - 4\Lambda})/2\Lambda$. For $\Lambda = 1/4$ and $M = 2\sqrt{2}/3$ all three zeros merge to a triple zero at $r = \sqrt{2}$. As long as μ_{RN} has three positive zeros the solution describes a charged black hole located inside a cosmological horizon. The horizon at the smallest value of r , a cosmological horizon according to our terminology, corresponds to the Cauchy horizon of the Reissner-Nordström black hole. The two possible double zeros at $r = r_{\pm}$ are the positions of extremal horizons of the black hole resp. cosmological type. Only the extremal black hole horizon has a limit for $\Lambda \rightarrow 0$.

²We call a horizon where r grows, while σ in Eq. (2) becomes positive a black hole horizon and a horizon, where r grows, while σ becomes negative, a cosmological horizon.

Apart from these general solutions of deSitter type there are some degenerate ones with r and W constant (and hence $N = U = 0$) and a space-time of the form $M_2 \times S^2$. In fact these solutions correspond to double zeros of μ_{SdS} or μ_{RN} , resp. to the SdS and RN fixed points (f.p.s) discussed in Section 3.8. From Eqs. (8,9) one finds that either $W^2 = 1$ (SdS) or $W = 0$ (RN) and $\dot{\kappa} = \kappa_s^2 - \kappa^2$ with

$$\kappa_s^2 = \sigma(1 - 2\Lambda r^2), \quad \text{and} \quad 2\Lambda r^2 = \begin{cases} 2 & \text{for } W^2 = 1, \\ 1 \pm \sqrt{1 - 4\Lambda} & \text{for } W = 0. \end{cases} \quad (19)$$

For $\kappa_s^2 > 0$ there are two solutions

$$\kappa(\tau) = \kappa_s \tanh(\kappa_s \tau), \quad e^{2\nu} = \cosh^2(\kappa_s \tau), \quad (20a)$$

$$\kappa(\tau) = \kappa_s \coth(\kappa_s \tau), \quad e^{2\nu} = \kappa_s^{-2} \sinh^2(\kappa_s \tau), \quad (20b)$$

with the line element $ds_2^2 = \sigma(e^{2\nu} dt^2 - r^2 d\tau^2)$ known as the Nariai solutions [9, 17]. For $\kappa_s^2 = -a^2 < 0$ one obtains

$$\kappa(\tau) = a \cot(a\tau), \quad e^{2\nu} = a^{-2} \sin^2(a\tau), \quad (20c)$$

and for $W = 0$ in the limit $\Lambda \rightarrow 1/4$, $\kappa_s \rightarrow 0$

$$\kappa(\tau) = \frac{1}{\tau}, \quad e^{2\nu} = \tau^2. \quad (20d)$$

Geometrically the first three are $SO(2,1)/SO(1,1)$, while the last one is flat space (in unusual coordinates).

3 Classification of Singular Points

Our aim is to give a complete classification of all solutions starting at an arbitrary regular point. We may integrate Eqs. (8) in both directions until either one of the dependent variables diverges for finite τ or $\tau \rightarrow \pm\infty$ where the solution may converge (to a finite or infinite value) or not. As the experience with a single Riccati equation shows we have to expect singularities for finite values of τ . In fact, a typical example is given by non-degenerate horizons, where the function κ has a simple pole. Natural candidates for the convergent case for $\tau \rightarrow \pm\infty$ are solutions running into one of the fixed points of Eqs. (8). The latter are easily found due to the simple structure of these equations. With exception of one case discussed in Section 3.5, the singularities occurring at finite values of τ turn out to be simple poles. These singularities can all be transformed into f.p.s by suitable changes of the independent and dependent variables. Furthermore all the f.p.s except the RN f.p. for $\Lambda = 1/4$ are hyperbolic ones. Using a technique known as ‘blow-up’ [20, 21] we are able to reduce also this exceptional case to the hyperbolic one.

Reducing the study of singular points to that of hyperbolic f.p.s has the advantage that the theory of dynamical systems provides powerful theorems about the local behaviour of solutions in their neighbourhood [18, 19, 20]. The

theorem of Hartman and Grobman about the existence of a stable (and unstable) manifold guarantees local existence and uniqueness of solutions. In order to parametrize the solutions by values determined at the f.p. itself, we rewrite the equations in the (non-autonomous and non-hyperbolic) form required by Prop. 12 in the appendix, whenever possible.

Most of the singularities are actually of geometrical origin. Since our independent variable τ parametrizes the space of orbits of the isometry group $G = R \times SO(3)$ of time-translations and rotations acting on the 4d space-time M_4 , we expect singularities at the fixed points of the group action. For rotations this is $r = 0$, the center of symmetry. It turns out that the point $r = 0$ can be reached for finite τ , if $|N| \rightarrow \infty$ (curvature singularity) or for $\tau \rightarrow \pm\infty$ with $N \rightarrow \mp 1$ (regular origin). Regular (non-degenerate) horizons are represented in the quotient space by their ‘bifurcation surface’ [22] consisting of f.p.s of the time-translations. The required vanishing of e^ν is possible only if κ diverges at some finite τ . In contrast, the degenerate horizons we find require $\tau \rightarrow \pm\infty$ and finite κ . A deSitter type singularity reached for finite τ occurs for $r \rightarrow \infty$ in the region $\sigma = -1$ due to the presence of the cosmological constant.

In the subsequent paragraphs we shall give a characterization of the solutions near the singular points discussed above, providing the basis for our classification theorems.

3.1 Regular Origin

As a first case there are solutions for $\sigma = +1$ with a regular origin $r \rightarrow 0$, $W^2 \rightarrow 1$, and $N \rightarrow \pm 1$ for $\tau \rightarrow \pm\infty$. These are f.p.s of Eqs. (8,9) with two dimensional stable manifolds. In order to characterize the corresponding one parameter families of solutions by a value b determined at the f.p., one rewrites Eqs. (8,9) in the form required by Prop. 12 as in [3]. Choosing $W \rightarrow +1$ without restriction one thus obtains unique solutions which are analytic in b and r with

$$W(r) = 1 - br^2 + O(r^4) , \quad (21a)$$

$$N^2(r) = 1 - \left(4b^2 + \frac{\Lambda}{3}\right)r^2 + O(r^4) . \quad (21b)$$

3.2 Reissner-Nordström Singularity at $r = 0$

The generic behaviour at $r = 0$ is singular in complete analogy to the case $\Lambda = 0$ [3], since the terms proportional to Λ may be neglected for $r \rightarrow 0$. The behaviour near $r = 0$ depends on the sign of σ . For $\sigma = +1$ one finds a Reissner-Nordström type singularity of the metric

$$W(r) = W_0 - \frac{W_0}{2(W_0^2 - 1)}r^2 + W_3r^3 + O(r^4) , \quad (22a)$$

$$N^2(r) = \frac{(W_0^2 - 1)^2}{r^2} - \frac{2M_0}{r} + O(1) , \quad (22b)$$

with three parameters M_0 , W_0 , and W_3 . As was shown in [16] the singularity may be described as a f.p. introducing a suitable set of dependent variables.

Here we shall give an alternative formulation using Schwarzschild coordinates and the dependent variables already introduced in Prop. 13 of [3].

$$r \frac{d}{dr} \bar{N} = r \left(\kappa \bar{N} - 2rU^2 \right) / \bar{N} , \quad (23a)$$

$$r \frac{d}{dr} \kappa = r \left(1 - 2\Lambda r^2 + 2U^2 - \kappa^2 \right) / \bar{N} , \quad (23b)$$

$$r \frac{d}{dr} W = r^2 U / \bar{N} , \quad (23c)$$

$$r \frac{d}{dr} U = r \left(\lambda - \kappa U \right) / \bar{N} , \quad (23d)$$

$$r \frac{d}{dr} \lambda = r \left(3W^2 - 2U^2 - 1 \right) U / \bar{N} , \quad (23e)$$

with $\bar{N} = rN$ and $\lambda = WT + NU$. These equations are precisely of the form required for the application of Prop. 12 implying the regularity of local solutions as a function of r obeying the constraints

$$\bar{N}(0) = \pm(W^2(0) - 1) , \quad \bar{N}(0)U(0) = W(0)(W^2(0) - 1) , \quad (24)$$

following from the finiteness of $\lambda(0)$ and the constraint Eq. (9). Geometrically this behaviour leads to a Reissner-Nordström type curvature singularity at which the space-time is geodesically incomplete.

3.3 Pseudo-Reissner-Nordström Singularity at $r = 0$

There is an analogous singularity for $\sigma = -1$ called pseudo-RN singularity in [16], with a rather different behaviour of the metric and of the YM potential W compared to the case $\sigma = +1$.

$$W(r) = W_0 \pm r + O(r^2) , \quad (25a)$$

$$N^2(r) = \frac{(W_0^2 - 1)^2}{r^2} \mp \frac{4W_0(W_0^2 - 1)}{r} + O(1) , \quad (25b)$$

The description as a f.p. was already given in [16], but here we give an alternative description, providing a unified treatment of the two different signs of W' at $r = 0$. We introduce the dependent variables $\bar{U} = U/N$, $\bar{T} = T/N$,

$$x = \frac{\bar{U}^2 - 1}{r} + \frac{2W\bar{U}\bar{T}^2}{W^2 - 1} , \quad \text{and} \quad y = \frac{\bar{T}^2 - 1}{r} . \quad (26)$$

Using Schwarzschild coordinates one finds

$$r \frac{d}{dr} W = r f_W , \quad f_W = \bar{U} , \quad (27a)$$

$$r \frac{d}{dr} \bar{U} = r f_U , \quad f_U = -\frac{W\bar{T}^2}{W^2 - 1} + 2z\bar{U} , \quad (27b)$$

$$r \frac{d}{dr} \bar{T} = r f_T , \quad f_T = \left[x + z - \frac{2W\bar{U}(\bar{T}^2 - 1)}{W^2 - 1} \right] \bar{T} , \quad (27c)$$

$$r \frac{d}{dr} x = -x - 2y + r f_x, \quad f_x = -2xy + 2 \left[2(x + y + 2z)W\bar{U} + \bar{U}^2 + \frac{(1 - \Lambda r^2)\bar{U}^2 + W^2(2\bar{U}^2 - \bar{T}^2 - 4\bar{U}^2\bar{T}^2)}{W^2 - 1} \right] \frac{\bar{T}^2}{W^2 - 1}, \quad (27d)$$

$$r \frac{d}{dr} y = 2x - 2y + r f_y, \quad f_y = 2(x + z)y + \left[\frac{1 - \Lambda r^2}{W^2 - 1} - 4W\bar{U}y \right] \frac{\bar{T}^2}{W^2 - 1}, \quad (27e)$$

with $z = [r(1 - \Lambda r^2)\bar{T}^2/(W^2 - 1)^2 - y]/2$. In their linearized form these equations have zero eigenvalues for W , $\bar{U}$, and $\bar{T}$, while x and y contribute with the complex conjugate eigenvalues $(-3 \pm \sqrt{15}i)/2$ representing divergent modes. Introducing the linear combinations (eigenmodes) $x_{\pm} = (1 \pm \sqrt{15}i)x - 4y$ we obtain

$$r \frac{d}{dr} x_{\pm} = \frac{-3 \pm i\sqrt{15}}{2} x_{\pm} + r f_{\pm}, \quad f_{\pm} = (1 \pm \sqrt{15}i)f_x - 4f_y, \quad (27f)$$

satisfying the requirements of Prop. 12. We thus obtain local solutions of Eqs. (27) with $x(0) = y(0) = 0$ depending analytically on $W(0) \neq \pm 1$, $\bar{U}(0)$, $\bar{T}(0)$, and r . Imposing Eqs. (26) as constraints we then obtain the conditions $\bar{U}^2(0) = \bar{T}^2(0) = 1$ and from the constraint Eq. (9) we find $\kappa/N = 1 + \bar{U}^2 - rz \rightarrow 2$ for $r \rightarrow 0$.

3.4 Schwarzschild Singularity at $r = 0$

The form of the RN type singularities requires that $W^2(0) \neq 1$, while for $W^2(0) = 1$ we obtain a Schwarzschild type behaviour ($N^2 \sim 1/r$) with the sign of the mass determined by σ . This case requires a different linearization, which we shall give subsequently. Assuming without restriction $W(0) = +1$, we introduce the dependent variables $\bar{W} = (W - 1)/r^2$, $\bar{U} = U/rN$, $k = \kappa/N$, $\bar{N} = \sigma r N^2$, and $y = (1 - 2\kappa/N + 2r^2\bar{U}^2)/r$, use the constraint Eq. (9) to express $y = \bar{N}^{-1}(1 - r^2(2 + r^2\bar{W})^2\bar{W}^2 - \Lambda r^2)$, and find

$$r \frac{d}{dr} \bar{N} = r f_N, \quad f_N = (y - 2r\bar{U}^2)\bar{N}, \quad (28a)$$

$$r \frac{d}{dr} \bar{W} = \bar{U} - 2\bar{W}, \quad (28b)$$

$$r \frac{d}{dr} \bar{U} = r f_U, \quad f_U = \frac{(1 + r^2\bar{W})(2 + r^2\bar{W})\bar{W}}{\bar{N}} + y\bar{U}. \quad (28c)$$

In order to bring Eq. (28b) into the form required by Prop. 12, we further introduce $\hat{W} = \bar{W} - \bar{U}/2$ with

$$r \frac{d}{dr} \hat{W} = -2\hat{W} - \frac{r f_U}{2}, \quad (28d)$$

and obtain solutions with $(\bar{W}, \bar{U}, \bar{N})$ tending to finite values $(-b, -2b, -2M)$ with $\sigma M < 0$. The corresponding asymptotic behaviour at $r = 0$ is

$$W = 1 - br^2 + O(r^4), \quad (29a)$$

$$\sigma N^2 = -\frac{2M}{r} + 1 + O(r) , \quad (29b)$$

$$\kappa = \frac{N}{2}(1 + O(r)) . \quad (29c)$$

The RN and S type singularities at $r = 0$ are different singular points with different asymptotic behaviour. The S case (for $\sigma = +1$) can, however, be seen as the limit $W_0 \rightarrow \pm 1$ of the more general RN case. In Fig. 1 we present numerical solutions for $\Lambda = 0.8$ starting at a regular origin with $b \sim 0.237$ as they approach the RN type singularity with $W_0 \approx -1$. N and κ grow like $1/\sqrt{r}$ and $U \approx 0$ as for the S type singularity in an interval that increases as $|W_0^2 - 1|$ decreases until eventually N starts to grow like $1/r$ whereas κ has a finite limit and U tends to $\pm W_0$. Thus in the limit $|W_0| \rightarrow 1$ the functions $W(r)$, $U(r)$, $N(r)$, and $\kappa(r)$ converge pointwise to those for the S type singularity.

3.5 Unbounded Oscillations at $r = 0$

The nature of the f.p. is rather different in the RN and pseudo-RN case. While for $\sigma = +1$ the f.p. is a simple attractor it is a partially repulsive (in the $\bar{U}$, $\bar{T}$ subsystem) focal point for $\sigma = -1$.

The RN singularity at the origin for $\sigma = +1$ is therefore generic (i.e. persistent for sufficiently small changes in the initial data and Λ), as is the deSitter asymptotics for $\sigma = -1$ described below. All other singular points discussed above and below are not generic because they require one or more divergent modes to be suppressed.

The generic behaviour for $\sigma = -1$ and r decreasing is somewhat more complicated. As described in [15, 16] this leads to an oscillatory behaviour of the solutions as they approach $r = 0$ with $\kappa \rightarrow -\infty$ and unbounded oscillating U and N . We introduce new dependent variables $n = N/\kappa$, $u = U/\kappa$, and $t = T/\kappa$ which are bounded due to Eq. (9) since κ diverges. In addition we introduce a new independent variable ρ with $d\rho = |\kappa|d\tau$ such that $\rho \rightarrow \infty$ as $\kappa \rightarrow -\infty$ (see the proof of Prop. 10 for details) and denote the derivative with respect to ρ by a prime. Using $z = 1/|\kappa|$ we find

$$r' = -rn , \quad (30a)$$

$$u' = (2u^2 - n + (2\Lambda r^2 - 1)z^2)u - Wzt , \quad (30b)$$

$$t' = (2u^2 + n - 1 + (2\Lambda r^2 - 1)z^2)t + 2Wzu , \quad (30c)$$

$$n' = (2u^2 + n - 2 + (2\Lambda r^2 - 1)z^2)n + 2u^2 , \quad (30d)$$

$$z' = (2u^2 - 1 + (2\Lambda r^2 - 1)z^2)z , \quad (30e)$$

and the constraint

$$2n = n^2 + 2u^2 + t^2 + z^2(\Lambda r^2 - 1) , \quad (30f)$$

with $|W|\sqrt{z} = \sqrt{z - rt}$ bounded. Neglecting terms that vanish as $z \rightarrow 0$ we obtain the asymptotic equations

$$u' = (2u^2 - n)u , \quad (31a)$$

$$t' = (2u^2 + n - 1)t , \quad (31b)$$

$$n' = (2u^2 + n - 2)n + 2u^2 , \quad (31c)$$

with the constraint

$$2n = n^2 + 2u^2 + t^2 , \quad (31d)$$

or, in terms of new dependent variables $\xi = n = N/\kappa$, $\eta = 2u^2/n = 2U^2/N\kappa$, and $\zeta = t^2/n = T^2/N\kappa$

$$\frac{\xi'}{\xi} = \xi - 2 + (\xi + 1)\eta = \xi(2 - \xi) - (\xi + 1)\zeta , \quad (32a)$$

$$\frac{\eta'}{\eta} = 2 - \eta + (\eta - 3)\xi = -(2 - \eta)^2 + (3 - \eta)\zeta , \quad (32b)$$

$$\frac{\zeta'}{\zeta} = \zeta - 2 + (4 - \xi - \zeta)\xi = 2 - \zeta - (\eta + \zeta)\eta , \quad (32c)$$

with the constraint

$$2 = \xi + \eta + \zeta . \quad (32d)$$

Fig. 2 shows four trajectories as they emerge from the focal point $(\xi, \eta, \zeta) = (1/2, 1, 1/2)$ corresponding to the pseudo-RN singularity and rapidly approach the bounding triangle consisting of the three separatrices $\xi = 0$, $\eta = 0$, and $\zeta = 0$ connecting the three hyperbolic fixed points $A = (2, 0, 0)$ corresponding to the S singularity, $B = (0, 0, 2)$ corresponding to a regular horizon described below, and $C = (0, 2, 0)$.

The behaviour referred to as ‘chaotic’ in [16] is due to this rapid approach to the boundary. The terms proportional to W in the full Eqs. (30) may become relevant when $\eta \ll z$ or $\zeta \ll z$. A solution may then either end with an S singularity or regular horizon or nearly miss them with a resulting substantial change in t or u . There is no such instability for $\xi \ll z$.

3.6 Regular Horizons

In the case of a regular, i.e. non-degenerate horizon we may use $s = 1/\kappa$ as the independent variable tending to zero. We thus replace Eqs. (8) by

$$s \frac{d}{ds} \tau = -\frac{\kappa}{\dot{\kappa}} = s (1 + s^2 F(r, U)) , \quad (33a)$$

$$s \frac{d}{ds} r = s (1 + s^2 F(r, U)) r N , \quad (33b)$$

$$s \frac{d}{ds} W = s (1 + s^2 F(r, U)) r U , \quad (33c)$$

$$s \frac{d}{ds} U = -U + s (1 + s^2 F(r, U)) (\sigma W T + N U) - s^2 F U , \quad (33d)$$

with

$$F(r, U) = \frac{1}{1 - s^2 f} , \quad f = \sigma (1 - 2\Lambda r^2) + 2U^2 , \quad (34)$$

while the function N can be computed from the constraint Eq. (9). These equations have precisely the form required in Prop. 12 with $U \rightarrow 0$ and τ, r, W tending to finite values τ_h, r_h, W_h .

In terms of the coordinate τ the behaviour near the horizon is (performing a shift in τ we arrange for $\tau_h = 0$)

$$r(\tau) = r_h \left(1 + N_1 \frac{\tau^2}{2} \right) + O(\tau^4), \quad (35a)$$

$$N(\tau) = N_1(\tau + O(\tau^3)) - W_1^2 \tau^3 + O(\tau^5), \quad (35b)$$

$$W(\tau) = W_h + r_h W_1 \frac{\tau^2}{2} + O(\tau^4), \quad (35c)$$

where

$$N_1 = \frac{\sigma}{2}(1 - \Lambda r^2 - T^2) \Big|_h, \quad W_1 = \frac{\sigma}{2} W T \Big|_h. \quad (36)$$

According to our definitions we have a black hole horizon if $\sigma N_1 > 0$ or a cosmological horizon if $\sigma N_1 < 0$. If N_1 vanishes the expansion of N starts with the cubic term with a corresponding change in the expansion of r . Geometrically this describes the situation that the horizon coincides with an equator, i.e. a maximum of r . The condition $\sigma N_1 > 0$ determines the domains $C_+ = \{1 - \Lambda r^2 - T^2 > 0\}$ in the (r, W) plane. In terms of r^2 and W^2 the boundary curves are (parts of) ellipses. In Fig. 3 these domains are shown for various values of Λ , for $\Lambda > 1/4$ they become disconnected.

Solutions with a regular origin can be seen as a limit $r_h \rightarrow 0$ of those with a regular horizon. In Fig. 4 we present solutions for $\Lambda \sim 0.3642$ starting at a regular origin with $b \sim 0.4296$ (i.e., near the point R_2 of Fig. 14) as they approach a regular horizon of decreasing radius. $N \approx \kappa \approx -1$ and $U/r \approx 2b$ as for the regular 3-sphere solution with 2 zeros of W in an interval that increases as $r_h \rightarrow 0$ until eventually κ starts to diverge whereas N and U tend to zero. Thus the black hole solutions converge to the one with a regular origin.

3.7 deSitter Asymptotics

For $\sigma = -1$ the radial variable r can tend to ∞ leading to deSitter asymptotics

$$W = W_0 + \frac{W_1}{r} + O(1/r^2), \quad (37a)$$

$$N^2 = \frac{\Lambda}{3} r^2 - 1 + \frac{2M_0}{r} + O(1/r^2). \quad (37b)$$

We use $s = 1/r$ as independent variable and introduce the dependent variables $\bar{U} = UN$, $\bar{N} = N/r$, and the ‘mass function’ Eq. (13).

$$s \frac{d}{ds} W = -s \frac{\bar{U}}{\bar{N}^2}, \quad (38a)$$

$$s \frac{d}{ds} \bar{U} = s \left(W(W^2 - 1) + 2s^3 \frac{\bar{U}^3}{\bar{N}^4} \right), \quad (38b)$$

$$s \frac{d}{ds} m = s \left(\frac{\bar{U}^2}{\bar{N}^2} - (W^2 - 1)^2 \right), \quad (38c)$$

with $\bar{N}^2 = \Lambda/3 - s^2 + 2s^3 m$. Thus all the requirements of Prop. 12 are fulfilled.

3.8 The Schwarzschild-deSitter and Reissner-Nordström Fixed Points

Besides the singular points reached at finite τ there are the f.p.s of Eqs. (8) with $(r, W, U, N, \kappa) \rightarrow (r_s, W_s, U_s, N_s, \kappa_s)$ as $\tau \rightarrow \infty$. One such f.p. is the regular origin $r_s = 0$ already discussed. For finite $r_s \neq 0$ this requires $U_s = N_s = 0$ and either $W_s^2 = 1$ or $W_s = 0$, which we denote as Schwarzschild-deSitter (SdS) resp. Reissner-Nordström (RN) f.p. The finite values of κ_s and r_s are determined by Eqs. (8d) and (9). In order to prove the local existence of solutions running into these f.p.s we will apply the usual linearization technique, using variables $\bar{r} = r/r_s - 1$, $\bar{W} = (W - W_s)/r$, and $\bar{\kappa} = \kappa - \kappa_s$.

The SdS f.p. requires $\sigma = -1$ in order to get $\kappa_s^2 > 0$ and we find $r_s = 1/\sqrt{\Lambda}$ and $\kappa_s = \pm 1$. From Eqs. (8) we obtain

$$\dot{N} = \kappa_s N + f_1, \quad (39a)$$

$$\dot{\bar{\kappa}} = -2\kappa_s(\bar{\kappa} + N) + f_2, \quad (39b)$$

$$\dot{\bar{W}} = U + f_3, \quad (39c)$$

$$\dot{U} = -2\bar{W} - \kappa_s U + f_4, \quad (39d)$$

and the constraint Eq. (9) yields $\bar{r} = N/\kappa_s + f_5$. The functions f_i , $i = 1, \dots, 5$ are analytic and at least quadratic in the dependent variables $\bar{W}$, U , N , and $\bar{\kappa}$. The eigenvalues of the linear system are $\kappa_s = \pm 1$, $-2\kappa_s = \mp 2$ for the N , κ part and $(-\kappa_s \pm \sqrt{7}i)/2$ in the $\bar{W}$, U sector. For $\kappa_s = 1$ there are three convergent modes for $\tau \rightarrow \infty$, while for $\kappa_s = -1$ there is only one convergent mode. For $\kappa_s = 1$ the function W performs damped oscillations of period $4\pi/\sqrt{7}$ around the limiting value $W_s = \pm 1$. For $\kappa_s = -1$ both divergent modes of the W , U subsystem have to be suppressed. However, the r.h.s. of Eqs. (39c,d) vanish for $\bar{W} = U = 0$ and thus the stable manifold of the f.p. contains only the SdS solution Eq. (17) with $W \equiv \pm 1$ and a double zero of μ_{SdS} . Similarly for $\kappa_s = +1$ the divergent mode of N has to be suppressed and thus $N(\tau) \approx \int_{\tau}^{\infty} 2U^2 e^{\tau-\tau'} d\tau' \geq 0$. Except for the Nariai solutions (20a,b) with $\bar{W} \equiv \bar{r} \equiv 0$, the f.p. can only be reached from $r < r_s$ with $N > 0$.

The RN f.p. with $W_s = 0$ leads to $r_s^2 = r_{\pm}^2 = (1 \pm \sqrt{1-4\Lambda})/2\Lambda$ and $\kappa_s^2 = \sigma(1-2\Lambda r_{\pm}^2)$. It requires $\Lambda \leq 1/4$ and the combinations $\sigma = +1$, r_- resp. $\sigma = -1$, r_+ to obtain $\kappa_s^2 \geq 0$. For the linearization we assume $\Lambda < 1/4$, i.e. $\kappa_s \neq 0$ and obtain

$$\dot{N} = \kappa_s N + f_1, \quad (40a)$$

$$\dot{\bar{\kappa}} = -4\sigma\Lambda r_s^2 \frac{N}{\kappa_s} - 2\kappa_s \bar{\kappa} + f_2, \quad (40b)$$

$$\dot{\bar{W}} = U + f_3, \quad (40c)$$

$$\dot{U} = -\sigma\bar{W} - \kappa_s U + f_4, \quad (40d)$$

where $\bar{r} = N/\kappa_s + f_5$ is again obtained from the constraint. The eigenvalues of the linear system are κ_s , $-2\kappa_s$ for N , κ and $(-\kappa_s \pm \sqrt{\kappa_s^2 - 4\sigma})/2$ for the $\bar{W}$, U system. Since $\kappa_s^2 < 1$ the latter eigenvalues are complex conjugate for $\sigma = +1$ and real for $\sigma = -1$. For $\sigma = +1$ there are three convergent modes for $\kappa_s > 0$

and only one convergent mode for $\kappa_s < 0$, whereas for $\sigma = -1$ there are always two convergent modes. The need to suppress the divergent modes for $\sigma = +1$ and $\kappa_s < 0$ again allows only the local (and thus global) solution $W \equiv 0$, i.e. the stable manifold contains only the RNdS solution. The period of the damped oscillations of W around zero in the case $\sigma = +1$, $\kappa_s > 0$ is $4\pi/\sqrt{4 - \kappa_s^2}$ and, as in the SdS case, the f.p. can only be reached from $r < r_s$ with $N > 0$, except for the Nariai solutions with $W \equiv \bar{r} \equiv 0$. For $\sigma = -1$, $\kappa_s > 0$ there are again the Nariai solutions and solutions with $W \neq 0$, $N > 0$, and $r < r_s$. For $\sigma = -1$, $\kappa_s < 0$ there is the RNdS solution and there are solutions with $W \neq 0$.

3.9 The Reissner-Nordström Fixed Point for $\Lambda = 1/4$

For $\Lambda = 1/4$ some of the eigenvalues of the linearized Eqs. (40) vanish and this degenerate case requires special consideration. First we introduce new dependent variables

$$\begin{aligned} y_W &\equiv \bar{W} = W/r, & y_U &\equiv \bar{U} = U + \kappa\bar{W}/2, \\ y_\kappa &\equiv \kappa, & y_r &\equiv \bar{r} = 1 - r^2/2, & y_N &\equiv \bar{N} = N + \bar{W}\bar{U}, \end{aligned} \quad (41)$$

and in addition

$$y_A \equiv A = \sigma\bar{W}^2 + \bar{U}^2. \quad (42)$$

Next we rewrite Eqs. (8) and the constraint Eq. (9) as

$$\dot{y}_i = F_i(y) + f_i(y), \quad i = W, U, \kappa, r, N, A, \quad (43a)$$

$$0 = F_c(y) + f_c(y), \quad (43b)$$

with the ‘leading’ contributions

$$\begin{pmatrix} F_W \\ F_U \end{pmatrix} = \begin{pmatrix} -\kappa/2 & 1 \\ -\sigma & -\kappa/2 \end{pmatrix} \begin{pmatrix} \bar{W} \\ \bar{U} \end{pmatrix}, \quad (44a)$$

$$F_\kappa = \sigma\bar{r} - \kappa^2, \quad (44b)$$

$$F_r = -2\bar{N}, \quad (44c)$$

$$F_N = -\frac{\sigma\bar{r}^2}{4}, \quad (44d)$$

$$F_A = -\kappa A, \quad (44e)$$

$$F_c = A - \kappa\bar{N} - \frac{\sigma\bar{r}^2}{4}, \quad (44f)$$

dominating the behaviour near the f.p., and with the ‘non-leading’ corrections

$$\begin{pmatrix} f_W \\ f_U \end{pmatrix} = \begin{pmatrix} -N & 0 \\ \bar{f}_U & N \end{pmatrix} \begin{pmatrix} \bar{W} \\ \bar{U} \end{pmatrix}, \quad \bar{f}_U = \frac{\sigma\bar{r}}{2} - \frac{\kappa^2}{4} + \sigma r^2 \bar{W}^2 + U^2 - \kappa N, \quad (45a)$$

$$f_\kappa = 2U^2, \quad (45b)$$

$$f_r = 2(\bar{W}\bar{U} + \bar{r}N), \quad (45c)$$

$$f_N = -\frac{\sigma\bar{r}^3}{2r^2} - \frac{N^2}{2} + \left(\bar{f}_U - \frac{\kappa^2}{4} - \frac{\sigma r^2 \bar{W}^2}{2}\right)\bar{W}^2, \quad (45d)$$

$$f_A = 2N(\bar{U}^2 - \sigma\bar{W}^2) + 2\bar{f}_U\bar{W}\bar{U}, \quad (45e)$$

$$f_c = -\frac{\sigma\bar{r}^3}{2r^2} + \frac{N^2}{2} + \left(\frac{\kappa^2}{4} - \frac{\sigma r^2 \bar{W}^2}{2}\right)\bar{W}^2. \quad (45f)$$

From the linearized form of these equations we get vanishing eigenvalues from κ , $\bar{r}$, $\bar{N}$, and A , whereas $\bar{W}$ and $\bar{U}$ contribute the eigenvalues $\pm i$ for $\sigma = +1$ or ± 1 for $\sigma = -1$. We therefore expect that (at least some of) the functions decrease with powers of τ instead of exponentially as for non-vanishing eigenvalues.

In order to analyze the behaviour of solutions tending to the f.p. as $\tau \rightarrow \infty$ we perform a ‘quasi-homogeneous blow-up’, following [21] with some minor variations. In terms of rescaled variables

$$\tilde{y}_i = \lambda^{-\alpha_i} y_i, \quad \alpha_\kappa = 1, \quad \alpha_W = \alpha_U = \alpha_r = 2, \quad \alpha_N = 3, \quad \alpha_A = 4, \quad (46)$$

we find

$$F_i(y) = \lambda^{\alpha_i+1} F_i(\tilde{y}), \quad f_i(y) = O(\lambda^{\alpha_i+2}), \quad \text{for } i = \kappa, r, N, A, \quad (47a)$$

$$F_c(y) = \lambda^4 F_c(\tilde{y}), \quad f_c(y) = O(\lambda^5). \quad (47b)$$

Furthermore the leading contributions F_i , $i = \kappa, r, N, A$ and F_c depend only on κ , $\bar{r}$, $\bar{N}$, and A . Hence they define a quasi-homogeneous dynamical system for $(\kappa, \bar{r}, \bar{N}, A)$ of type $\alpha = (1, 2, 3, 4)$ and degree 2 in the sense of [21], whereas the non-leading corrections yield at least one additional power of λ upon rescaling. The non-leading correction terms, f_i depend on $\bar{W}$ and $\bar{U}$ individually, not only on the combination A . One can easily find bounds for $|\bar{W}|$ and $|\bar{U}|$: for $\sigma = +1$ they are bounded by $\sqrt{\tilde{A}}$, whereas for $\sigma = -1$ with eigenvalues ± 1 they are bounded by $\exp(-\delta\tau)$ for any $\delta < 1$ due to the special form of Eqs. (44a,45a).

Next we introduce $\mu_i = M/\alpha_i$ with some M such that $\mu_i \geq 2$, new dependent variables λ and $\tilde{y}$

$$\lambda^M = \sum_i \gamma_i |y_i|^{\mu_i}, \quad \tilde{y}_i = \lambda^{-\alpha_i} y_i, \quad i = \kappa, r, N, A, \quad (48)$$

with some constants $\gamma_i \geq 0$ such that $\sum_i \gamma_i > 0$, and a new independent variable t with $dt = \lambda d\tau$. For $\gamma_i > 0$ this is a ‘polar’ blow-up whereas e.g., $\gamma = (1, 0, 0, 0)$ yields a ‘directional’ blow-up in the κ -direction. Neglecting for a moment the non-leading correction terms we obtain

$$\frac{1}{\lambda} \frac{d\lambda}{dt} = \sum_i \gamma_i |\tilde{y}_i|^{\mu_i} \frac{F_i(\tilde{y})}{\alpha_i \tilde{y}_i}, \quad (49a)$$

$$\frac{d\tilde{y}_i}{dt} = F_i(\tilde{y}) - \alpha_i \tilde{y}_i \frac{1}{\lambda} \frac{d\lambda}{dt}, \quad (49b)$$

i.e., an $\tilde{y}$ -dependent ‘radial’ flow with the f.p. $\lambda = 0$ and a λ -independent ‘tangential’ flow with f.p.s determined by the conditions

$$\alpha_i \tilde{y}_i F_j(\tilde{y}) = \alpha_j \tilde{y}_j F_i(\tilde{y}), \quad i, j = \kappa, r, N, A. \quad (50)$$

There are three pairs of tangential f.p.s $P_{i\pm}$, $i = 1, 2, 3$ with $\tilde{\kappa} \gtrless 0$ and

$$P_{1\pm}: \quad \tilde{r} = \tilde{N} = \tilde{A} = 0, \quad (51a)$$

$$P_{2\pm}: \quad \tilde{r} = \frac{3\sigma}{4} \tilde{\kappa}^2, \quad \tilde{N} = \frac{3\sigma}{16} \tilde{\kappa}^3, \quad \tilde{A} = \frac{21\sigma}{64} \tilde{\kappa}^4, \quad (51b)$$

$$P_{3\pm}: \quad \tilde{r} = \frac{4\sigma}{3} \tilde{\kappa}^2, \quad \tilde{N} = -\frac{4\sigma}{9} \tilde{\kappa}^3, \quad \tilde{A} = 0. \quad (51c)$$

Inserting these values into Eq. (49a) yields $d\lambda/dt = -\tilde{\kappa}\lambda$ at $P_{1\pm}$, $-\tilde{\kappa}\lambda/4$ at $P_{2\pm}$, and $+\tilde{\kappa}\lambda/3$ at $P_{3\pm}$. Thus P_{1+} , P_{2+} , and P_{3-} are attractors for the radial flow, whereas P_{1-} , P_{2-} , and P_{3+} are repulsive. On the 2-dimensional surface $\tilde{y}$ subject to the constraint $F_c(\tilde{y}) = 0$ the f.p.s P_{1-} and P_{3+} are attractors for the tangential flow, P_{1+} and P_{3-} are repulsive, and $P_{2\pm}$ are saddle points. Furthermore there are trajectories along the curve $\tilde{N}^2 = \sigma\tilde{r}^3/12$ connecting all six f.p.s as shown in Fig. 5: separatrices from P_{1+} and P_{3-} to P_{2+} , separatrices from P_{2-} to P_{1-} and P_{3+} , as well as trajectories from P_{1+} to P_{3+} and from P_{3-} to P_{1-} . This leaves no room for limit or separatrix cycles, therefore according to the Poincaré-Bendixson theory the f.p.s are the only limit points of the tangential flow.

With this blow-up we have transformed the degenerate f.p. $y = 0$ into a set of hyperbolic f.p.s $(\lambda, \tilde{y}) = (0, P_{i\pm})$ with one dimensional stable manifolds for $(0, P_{1+})$, $(0, P_{2-})$, $(0, P_{3-})$ and two dimensional stable manifolds for $(0, P_{1-})$, $(0, P_{2+})$, $(0, P_{3+})$. The non-leading correction terms, f_i in Eqs. (43) yield $O(\lambda)$ corrections to the r.h.s. of Eqs. (49) and thus preserve the position and character of the six f.p.s. The structural stability of the system (43) guarantees that the flow pattern is not changed by the non-leading corrections. Moreover the stable manifolds for $(0, P_{1-})$, $(0, P_{2-})$, and $(0, P_{3+})$ are purely tangential at $\lambda = 0$ and therefore excluded for trajectories tending to the f.p. $y = 0$ as $\tau \rightarrow \infty$.

We have chosen $\bar{W}$, $\bar{U}$, and $\bar{N}$ such that $1/\lambda - c_i\tilde{\kappa}\tau$ has a finite limit on each of the stable manifolds with $c_1 = 1$ for P_{1+} , $c_2 = 1/4$ for P_{2+} , and $c_3 = -1/3$ for P_{3-} . Consequently we choose yet another independent variable $s = \lambda = 1/\tau$ as well as dependent variables $\tilde{y}$ as in Eq. (46) to obtain

$$s \frac{d}{ds} \begin{pmatrix} \tilde{W} \\ \tilde{U} \end{pmatrix} = \begin{pmatrix} \tilde{\kappa}/2 - 2 & -1/s \\ \sigma/s & \tilde{\kappa}/2 - 2 \end{pmatrix} \begin{pmatrix} \tilde{W} \\ \tilde{U} \end{pmatrix} - s \begin{pmatrix} g_W & 0 \\ \bar{g}_U & g_U \end{pmatrix} \begin{pmatrix} \tilde{W} \\ \tilde{U} \end{pmatrix},$$

$$g_W = -\frac{N}{s^2}, \quad g_U = \frac{N}{s^2}, \quad \bar{g}_U = \frac{\bar{f}_U}{s^2}, \quad (52a)$$

$$s \frac{d}{ds} \tilde{\kappa} = \tilde{\kappa}^2 - \tilde{\kappa} - \sigma\tilde{r} - sg_\kappa, \quad g_\kappa = \frac{f_\kappa}{s^3}, \quad (52b)$$

$$s \frac{d}{ds} \tilde{r} = 2(\tilde{N} - \tilde{r}) - sg_r, \quad g_r = \frac{f_r}{s^4}, \quad (52c)$$

$$s \frac{d}{ds} \tilde{N} = \frac{\sigma\tilde{r}^2}{4} - 3\tilde{N} - sg_N, \quad g_N = \frac{f_N}{s^5}, \quad (52d)$$

$$s \frac{d}{ds} \tilde{A} = (\tilde{\kappa} - 4)\tilde{A} - sg_A, \quad g_A = \frac{f_A}{s^6}, \quad (52e)$$

as well as the constraint

$$0 = \tilde{A} - \tilde{\kappa}\tilde{N} - \frac{\sigma\tilde{r}^2}{4} + sg_c, \quad g_c = \frac{f_c}{s^5}, \quad (52f)$$

with g 's that are analytic in s and the rescaled functions. The requirement that the r.h.s. of Eqs. (52b-e) vanish at $s = 0$ yields the conditions Eqs. (51) for the f.p.s with $\tilde{\kappa}(0) = 1$ for P_{1+} , $\tilde{\kappa}(0) = 4$ for P_{2+} , and $\tilde{\kappa}(0) = -3$ for P_{3-} . These three different solutions are to be expected as the limit $\Lambda \rightarrow 1/4$ of the Nariai solution (20b), and the RN f.p.s with the combinations r_+ , $\kappa > 0$ and

r_- , $\kappa < 0$. In terms of shifted variables $\hat{r} = \tilde{r} - \tilde{r}(0)$ etc., that vanish at $s = 0$ we thus obtain the linearized equations

$$s \frac{d}{ds} \begin{pmatrix} \hat{r} \\ \hat{N} \\ \hat{\kappa} \\ \hat{A} \end{pmatrix} = \begin{pmatrix} -2 & 2 & 0 & 0 \\ \sigma \frac{\tilde{r}(0)}{2} & -3 & 0 & 0 \\ -\sigma & 0 & 2\tilde{\kappa}(0) - 1 & 0 \\ 0 & 0 & \tilde{A}(0) & \tilde{\kappa}(0) - 4 \end{pmatrix} \begin{pmatrix} \hat{r} \\ \hat{N} \\ \hat{\kappa} \\ \hat{A} \end{pmatrix}, \quad (53)$$

with eigenvalues $+1$ (corresponding to a shift in τ), $\tilde{\kappa}(0) - 4$ (due to the constraint), as well as $-2, -3$ for P_{1+} , $+7, -6$ for P_{2+} , and $-6, -7$ for P_{3-} .

We still need an additional equation for the YM functions $\tilde{W}$ and/or $\tilde{U}$. For $\sigma = +1$ the solutions with $\tilde{\kappa}(0) < 4$ yield $W \equiv 0$ whereas $\tilde{\kappa}(0) = 4$ requires $W \neq 0$. Motivated by the eigenvalues $\pm i$, we introduce in the latter case $\tilde{W} = \sqrt{\tilde{A}} \sin(\tau + \theta)$ and $\tilde{U} = \sqrt{\tilde{A}} \cos(\tau + \theta)$. The equation

$$s \frac{d}{ds} \theta = s((g_U - g_W) \cos(\tau + \theta) + \bar{g}_U \sin(\tau + \theta)) \sin(\tau + \theta), \quad (54)$$

implies that the phase θ has a finite limit as $s \rightarrow 0$.

For $\sigma = -1$ with the eigenvalues ± 1 we introduce $\tilde{W}_\pm = \tilde{W} \pm \tilde{U}$ and replace Eq. (52f) by

$$s \frac{d}{ds} \tilde{W}_\pm = \left(\mp \frac{1}{s} + \frac{\tilde{\kappa} - 4}{2} \right) \tilde{W}_\pm - s(g_\pm \tilde{W}_\pm + \bar{g}_\pm \tilde{W}_\mp), \quad (55)$$

where $g_\pm$ and $\bar{g}_\pm$ are linear combinations of g_W , g_U , and $\bar{g}_U$. The divergent mode $\tilde{W}_+$ has to be suppressed and consequently $\tilde{W}_+$ and $\tilde{W}_-$ decrease exponentially, implying that $\tilde{A} = \tilde{W}_+ \tilde{W}_-$ vanishes at $s = 0$ and thus excluding the solution with $\tilde{\kappa}(0) = 4$. As for $\Lambda < 1/4$, there are solutions with $\kappa > 0$ and $\kappa < 0$ with both $W \equiv 0$ and $W \neq 0$.

4 Classification of Solutions

Starting from any regular point $\tau = \tau_0$ of the Eqs. (8) we may integrate these equations to both sides until we meet a singular point. In case $\int r d\tau$ is finite the corresponding space-time is geodesically incomplete, otherwise it is complete in the radial direction. This does not necessarily mean completeness in all directions as the example of the extremal RN solution demonstrates. A general recipe how to glue together incomplete pieces of space-time to a geodesically complete manifold is described in [23].

In the preceding section we have already discussed the possible singular points we may encounter integrating Eqs. (8) and described the local behaviour in their neighbourhood. The missing step is a classification of the singular points reachable from some regular point integrating the equations to both sides. Instead of integrating towards decreasing τ we may as well reverse the sign of N , κ , and U and always integrate towards increasing τ .

The global classification is different for the two possibilities $\sigma = \pm 1$, i.e. for regions with space-like or time-like radial direction. Here we start with some general properties equally valid for both cases.

Proposition 1: Consider a solution of Eqs. (8,9).

- i) If $N(\tau_0) < 0$ for some τ_0 then $N(\tau) < 0$ for all $\tau > \tau_0$.
- ii) r has no minima and at most one maximum.
- iii) r has a limit r_1 as $\tau \rightarrow \tau_1$ with $\ln r_1$ finite if and only if $\int^{\tau_1} N d\tau$ is finite.
- iv) If $N(\tau)$ and $\kappa(\tau)$ are finite at some τ then $W(\tau)$ and $U(\tau)$ are finite.
- v) If $N < 0$, $\kappa < 0$, and $\kappa + N$ is unbounded, then $\kappa + N \rightarrow -\infty$ at some finite τ_1 .

Let us define $\zeta = (\kappa - N)/(\kappa + N)$ and $\gamma = (\tau - \tau_1)(\kappa + N)$. If ζ is constant then $\gamma = 4/(1 + 3\zeta^2) + O((\tau - \tau_1)^2)$. If ζ is restricted to an interval then γ is some (τ dependent) average of this expression over that interval, in the most general case $1 \leq \gamma \leq 4$ up to terms $O((\tau - \tau_1)^2)$.

Proof:

- i) This is an immediate consequence of Eq. (8c).
- ii) This is an immediate consequence of the previous point and Eq. (8a).
- iii) Property ii) implies that r is monotonic for sufficiently large τ and therefore has a limit; the finiteness of $\ln r$ is a consequence of Eq. (8a).
- iv) Due to the previous point $\ln r$ has a finite limit and U^2 is integrable due to Eq. (8c). The Schwarz inequality then implies that $|U|$ and thus $r|U|$ are integrable, i.e. W is bounded. The boundedness of U finally follows from Lemma 13 in the appendix applied to Eq. (8e).
- v) From Eqs. (8c,d) we obtain

$$\begin{aligned} (\kappa + N)' &= \sigma(1 - 2\Lambda r^2) - \frac{1}{4}(\kappa + N)^2 - \frac{3}{4}(\kappa - N)^2 \\ &\leq \sigma(1 - 2\Lambda r^2) - \frac{1}{4}(\kappa + N)^2, \end{aligned} \quad (56)$$

with $1 - 2\Lambda r^2$ bounded since $N < 0$. Therefore Lemma 14 implies that $\kappa + N$ diverges for some finite τ_1 and we assume $\tau_1 = 0$ without restriction. Introducing $\eta = 1/(\tau(\kappa + N))$ we find

$$\eta(\tau) = -\frac{1}{4\tau} \int_{\tau}^0 (1 + 3\zeta^2) d\tau' + \sigma\tau \int_{\tau}^0 \left(\frac{\tau'}{\tau}\right)^2 (1 - 2\Lambda r^2) \eta^2 d\tau'. \quad (57)$$

Since $\eta \leq 1 + O(\tau^2)$ due to Lemma 14, the second integral on the r.h.s. is $O(\tau^2)$, whereas the first integral yields $1/\gamma$ as the average value of $(1 + 3\zeta^2)/4$. ■

According to Prop. 1 we can choose a suitable $\bar{\tau}_0 \geq \tau_0$ such that N has a definite sign for $\tau \geq \bar{\tau}_0$. We have to consider the cases that either N and κ are bounded for all $\tau \geq \bar{\tau}_0$ or that N and/or κ are unbounded.

For N and κ bounded we have the result:

Proposition 2: Suppose a solution of Eqs. (8,9) with N , κ , W , U , and $\ln r$ bounded for $\tau \geq \tau_0$, then the solution tends to one of the f.p.s of Eqs. (8) as $\tau \rightarrow \infty$:

- i) for $\sigma = -1$, the SdS f.p. described by Eqs. (39);
- ii) for $\sigma = \pm 1$, the RN f.p. described by Eqs. (40).

Proof: Since $\ln r$ is bounded, $|N|$ is integrable and therefore U^2 is integrable due to Eq. (8c). Integrating

$$(U^3)' = 3\left(\sigma WT + (N - \kappa)U\right)U^2, \quad (58)$$

derived from Eq. (8e), we see that U has a limit, which must be zero since otherwise W would diverge. Lemma 14 applied to Eq. (8d) implies that κ has a limit. From Eqs. (14) then follows that the ‘energy’ E is bounded and $|\dot{E}|$ is integrable, thus E has a limit. This implies that N has a limit which must be zero since otherwise $\ln r$ would diverge, and that W has a finite limit which must be one of the f.p.s of Eq. (8e), $W \rightarrow \pm 1$ for the SdS f.p. or $W \rightarrow 0$ for the RN f.p. ■

For N bounded and κ unbounded we have the result:

Proposition 3: Suppose a solution of Eqs. (8,9) with N bounded and κ unbounded, then the solution has a regular horizon at some finite τ_1 as described in Eqs. (35,36).

Proof: First we show that τ_1 must be finite. For $\sigma = +1$ we use Eq. (56), for $\sigma = -1$ we use

$$(\kappa + N)' = 1 - 4U^2 - 2T^2 - \kappa^2 + 5\kappa N - 3N^2 \leq 1 + 7N(\kappa + N) - (\kappa + N)^2. \quad (59)$$

In both cases Lemma 15 implies that $\kappa + N$ and thus κ diverges to $-\infty$ for some finite τ_1 , thus $\ln r$ has a finite limit. We assume again $\tau_1 = 0$ without restriction.

Next, for $\sigma = +1$ Eq. (11) implies that U^2 is integrable, since otherwise N would be unbounded from below. Therefore $|U|$ is integrable due to the Schwarz inequality and thus W has a finite limit. The constraint Eq. (9) then implies that U^2/κ is bounded. For $\sigma = -1$ the constraint implies that U^2/κ is bounded.

For $\sigma = \pm 1$ we use $s = 1/\kappa$ as in Eqs. (33,34) with $d\tau = (1 + s^2 F) ds$ and find that sf and hence sF is bounded. Therefore $\int_0^0 |U| d\tau < \infty$ and thus W has a finite limit. We now turn to Eq. (33d) for U and obtain

$$\frac{d}{ds} \left(se^G \frac{U}{r} \right) = \sigma se^G (1 + s^2 F) \frac{W(W^2 - 1)}{r^2}, \quad (60)$$

with $G(s) = -\int_s^0 s'^2 F ds'$ bounded, yielding

$$e^G \frac{U}{r} = \frac{c}{s} - \sigma \int_s^0 \frac{s'}{s} e^G (1 + s'^2 F) \frac{W(W^2 - 1)}{r^2} ds'. \quad (61)$$

The constant c must vanish, otherwise sU^2 would not be bounded. This implies $U \rightarrow 0$ for $s \rightarrow 0$. From the constraint we finally obtain $N(0) = 0$, thus the solution has a horizon. ■

4.1 Solutions with $\sigma = +1$

To begin with we state some properties of the solutions required for their global classification. Most of these are trivial generalizations of results already proved for $\Lambda = 0$ in [3], but we repeat their proofs for completeness.

Proposition 4: *Consider a solution of Eqs. (8,9) with $\sigma = +1$.*

- i) *The function W can have neither maxima if $W > 1$ or $0 > W > -1$ nor minima if $W < -1$ or $0 < W < 1$.*
- ii) *N is bounded from above and if $N(\tau_0) < 1$ for some τ_0 then $N(\tau) < 1$ for all $\tau > \tau_0$.*
- iii) *r is bounded from above.*
- iv) *$\kappa + N$ is bounded from above and if $(\kappa + N)(\tau_0) < 2$ for some τ_0 then $(\kappa + N)(\tau) < 2$ for all $\tau > \tau_0$.*
- v) *If $W^2(\tau_0) > 1$ and $WU(\tau_0) \geq 0$ for some τ_0 and κ is bounded, then $N(\tau_1) < -1$ for some finite $\tau_1 > \tau_0$.*
- vi) *If $N(\tau_0) < -1$ for some τ_0 and κ is bounded, then $N(\tau)$ diverges to $-\infty$ for some finite $\tau_1 > \tau_0$.*

Proof:

- i) This is an immediate consequence of Eqs. (8b,e).
- ii) This follows from Eq. (11).
- iii) In the following argument we can always assume $\Lambda r^2 \geq 3$ and $N \geq 0$ since otherwise there is nothing to prove. Thus $\dot{N} \leq -1$ due to Eq. (11) and therefore $N = 0$ for some finite τ_1 where r has a finite maximum.
- iv) Eq. (56) implies $(\kappa + N) \leq 0$ for $\kappa + N \geq 2$.
- v) In the following argument we can always assume $N \geq -1$ since otherwise there is nothing to prove. Let $A > 1$ be a constant such that $\kappa < 2A$. First we note that W^2 monotonically increases and $WU > 0$ for $\tau > \tau_0$. From $\ddot{W} = W(W^2 - 1) + (2N - \kappa)\dot{W}$ we obtain a positive lower bound for $|\dot{W}|$ and thus there exists a $\tilde{\tau}_0 \geq \tau_0$ such that $|\dot{W}|(\tilde{\tau}_0) > A$. Using the Schwarz inequality we find

$$|TU|' = 2|W|(2T^2 + U^2) - \kappa|TU| \geq (\sqrt{8} - 2)|TU|. \quad (62)$$

Therefore $|TU|$ and, using again the Schwarz inequality, $2T^2 + U^2$ grow exponentially for $\tau > \tilde{\tau}_0$. Eq. (11) then implies that $N < -1$ for some finite τ_1 .

vi) This is a consequence of Eq. (11) and Lemma 14. ■

In addition there are new features compared to the case $\Lambda = 0$, which are mainly due to the possibility that the variable κ may become negative.

Proposition 5: *Suppose a solution of Eqs. (8,9) for $\sigma = +1$ with N and κ bounded for all $\tau \geq \tau_0$, then the solution runs into*

- i) *the RN f.p. described by Eqs. (40), if $r \rightarrow r_1 > 0$ (with $\kappa_s < 0$ only possible if $W \equiv 0$);*
- ii) *a regular origin as described by Eqs. (21), if $r \rightarrow 0$.*

Proof: First we note that the limit r_1 of r cannot exceed $1/\sqrt{\Lambda}$ since otherwise N would diverge due to Eq. (11). Moreover the ‘mass function’ Eqs. (13) is monotonic and bounded and therefore has a limit. Furthermore W is bounded due to Prop. 4.

- i) If $r_1 > 0$ then U is bounded due to the constraint Eq. (9) and Prop. 2 implies that the solution runs into the RN f.p., as discussed in Section 3. For reasons given there the case $\kappa_s < 0$ is only possible if $W \equiv 0$.
- ii) For $r_1 = 0$ we first of all choose a $\tilde{\tau}_0 \geq \tau_0$ such that $N < 0$ and $\Lambda r^2 < 1/2$ for all $\tau \geq \tilde{\tau}_0$ and restrict the following arguments to this subinterval.

Eq. (56) implies that $\kappa + N \geq -2$ since otherwise $\kappa + N$ would diverge. From Eq. (8d) we see that if $\kappa > -\sqrt{1 - 2\Lambda r^2}$ for some $\tilde{\tau}_0$, then κ will increase towards zero and stay positive afterwards. This is, however, impossible since then $(rN)' = \kappa rN - 2rU^2 \leq 0$ and rN would not converge to zero as required. We thus obtain the bounds $-2 \leq \kappa \leq -\sqrt{1 - 2\Lambda r^2}$.

We then show that W cannot have zeros for $r < 1/2$. Assume there were such a zero at $\tilde{\tau}_0$. This would imply $T^2 > 9/4$ for $\tau \geq \tilde{\tau}_0$ as long as $|W| < 1/2$. Thus $|W| = 1/2$ for some $\tau = \tilde{\tau}_0 + \Delta$ since otherwise N would diverge due to Eq. (11). We then estimate, repeatedly using the Schwarz inequality

$$\begin{aligned} N(\tilde{\tau}_0 + \Delta) &\leq N(\tilde{\tau}_0) + \int_{\tilde{\tau}_0}^{\tilde{\tau}_0 + \Delta} \frac{1}{2} \left(1 - \frac{2\dot{W}^2 + (W^2 - 1)^2}{r^2} \right) d\tau \\ &\leq -\frac{5\Delta}{8} - \frac{1}{\Delta} < -1, \end{aligned} \tag{63}$$

and thus N would diverge due to Prop. 4.

Since the extrema of W are separated by zeros, W is monotonic for sufficiently large τ and therefore has a limit. This limit can only be ± 1 since otherwise N would diverge due to Eq. (11).

Next we estimate (using $TUW \leq 0$)

$$|TU|' = -2|W|(T^2 + 2U^2) - \kappa|TU| \leq -(\sqrt{8}|W| - 2)|TU|, \tag{64}$$

i.e., $|TU|$ decreases exponentially in τ . Eq. (64) then implies that T and U both must decrease exponentially since TU cannot change sign.

Applying Lemma 14 to Eqs. (8d) and (11) we finally obtain $\kappa \rightarrow -1$ and $N \rightarrow 1$ for $\tau \rightarrow \infty$. Thus the solution tends to a regular origin. ■

Proposition 6: *Suppose a solution of Eqs. (8,9) for $\sigma = +1$ with N unbounded, then the solution has an origin $r = 0$ at some finite τ_1 with*

- i) *the S type singularity as described in Eqs. (28,29), if κ is unbounded;*
- ii) *the RN type singularity as described in Eqs. (22,23), if κ is bounded.*

Proof: In view of Prop. 4 N must be unbounded from below. Integrating Eq. (11) shows that $N \rightarrow -\infty$ for some finite τ_1 , and we assume $\tau_1 = 0$ without restriction. In the following we will consider sufficiently small $|\tau|$ such that $N(\bar{\tau}_0) < -1$. In order to proceed we distinguish the cases that κ is unbounded or bounded from below.

- i) For κ unbounded from below we first note that $\kappa + N$ is unbounded from below and thus $\kappa + N \rightarrow -\infty$ for $\tau \rightarrow 0$ due to Eq. (56).

From the equation

$$\begin{aligned} (2N - \kappa)' &= (N - \kappa)(2N - \kappa) + \frac{1}{2}(3 - 3N^2 - 2U^2 - 5T^2 - \Lambda r^2) \\ &< (N - \kappa)(2N - \kappa) , \end{aligned} \quad (65)$$

we then see that if $(2N - \kappa)(\bar{\tau}_0) \leq 0$ for some $\bar{\tau}_0$, then $2N - \kappa \leq 0$ for all $\tau \geq \bar{\tau}_0$.

First assume $2N - \kappa > 0$ for all τ and therefore $\kappa \rightarrow -\infty$. Now consider the equation

$$\begin{aligned} \left(\frac{\kappa}{N}\right)' &= (\kappa - 2N)\left(2 + \frac{T^2 + \Lambda r^2}{N^2}\right) + \frac{3T^2 + \Lambda r^2}{N} + 3N - \frac{1}{N} \frac{\kappa}{N} \\ &< 3N + \frac{1}{|N|} \frac{\kappa}{N} . \end{aligned} \quad (66)$$

With $A(\tau) = \int_{\tau}^0 d\tau' / |N| = O(\tau)$ we then find $(e^A \kappa / N)' < 3Ne^A < 0$ and therefore κ/N , which is non-negative, has a finite limit. Lemma 14 then implies $\int_{\tau}^0 \kappa d\tau' = -\infty$ and therefore $\int_{\tau}^0 N d\tau' = -\infty$. This is, however, impossible since κ/N would then become negative due to Eq. (66).

Therefore $2N - \kappa \leq 0$ for $|\tau|$ sufficiently small. Next we want to show that $\kappa < 0$ for $|\tau|$ sufficiently small. For $r \rightarrow 0$ this follows from Eq. (8d). For $r \rightarrow r_1 > 0$ we first assume that W is bounded. Lemma 13 applied to the equation

$$(rU)' = W(W^2 - 1) + (2N - \kappa)rU , \quad (67)$$

then implies that U is bounded and the constraint Eq. (9) implies $\kappa < 0$. Then we assume that W is unbounded and therefore, due to Prop. 4,

$WU > 0$ and $|W| \rightarrow \infty$. This requires that U is unbounded and the equation

$$\left(\frac{U}{r^2}\right)' = \frac{W(W^2 - 1)}{r^3} - (\kappa + N)\frac{U}{r^2}, \quad (68)$$

then implies that U/r^2 is monotonic and thus U diverges. Eq. (8d) then again implies $\kappa < 0$.

Now consider

$$(r\kappa + rN)' = r\kappa(2N - \kappa) + r(1 - 2\Lambda r^2). \quad (69)$$

The r.h.s. is bounded from below and therefore $r\kappa + rN$, which cannot be positive, has a finite limit. This implies that $r \rightarrow 0$ and that $r\kappa$ and rN are bounded. From the equation

$$(r^2N)' = (r\kappa + rN)rN - 2r^2U^2, \quad (70)$$

then follows that r^2U^2 is integrable and, using the Schwarz inequality, that W has a finite limit W_1 . Eq. (67) finally implies that $\dot{W} \equiv rU$ has a finite limit $\dot{W}_1$.

From Eqs. (13b) we obtain

$$m(\tau) = m(\tau_0) - \int_{r(\tau)}^{r(\tau_0)} \frac{2\dot{W}^2 + (W^2 - 1)^2}{2r^2} dr', \quad (71)$$

and therefore $\lim_{\tau \rightarrow 0} rm = -\dot{W}_1^2 - (W_1^2 - 1)^2/2$. Thus $r\kappa$ and rN have finite limits k_1 resp. n_1 satisfying $n_1^2 = 2\dot{W}_1 + (W_1^2 - 1)^2$ and, due to the constraint Eq. (9), $2k_1n_1 = n_1^2 + 2\dot{W}_1 - (W_1^2 - 1)^2 = 4\dot{W}_1^2$.

We want to show that $n_1 = 0$. Assuming $n_1 < 0$, we find $2n_1 - k_1 < 0$ and $r(\tau) = n_1\tau + O(\tau^2)$. This implies $\int_{\tau}^0 (2N - \kappa)d\tau' = -\infty$ and Lemma 13 applied to Eq. (67) yields $k_1 = \dot{W}_1 = 0$. As a consequence $\int_{\tau}^0 \kappa d\tau$ is finite and Eq. (8d) implies that κ is bounded from below. This is a contradiction, therefore $n_1 = k_1 = \dot{W}_1 = 0$ and $W_1^2 = 1$, and we may assume $W_1 = +1$ without restriction.

From the equation

$$(2\kappa - N)' = -(\kappa + N)(2\kappa - N) + 6U^2 + 2(1 - \Lambda r^2) > -(\kappa + N)(2\kappa - N), \quad (72)$$

we see that if $(2\kappa - N)(\bar{\tau}_0) > 0$ for some $\bar{\tau}_0$, then $2\kappa - N > 0$ for all $\bar{\tau}_0 \leq \tau < 0$. This is, however, impossible because then $(2r\kappa - rN) \geq 0$ and $2r\kappa - rN$ could not converge to zero as required. Therefore $2\kappa - N \leq 0$ for all sufficiently small $|\tau|$.

We can now rewrite Eq. (66) as

$$\left(\frac{\kappa}{N}\right)' = (2\kappa - N) + \frac{T^2}{N} \left(\frac{\kappa}{N} + 1\right) - \frac{\Lambda r^2}{N} + \frac{\Lambda r^2 - 1}{N} \frac{\kappa}{N}. \quad (73)$$

The first two terms on the r.h.s. are both negative and the last two terms are bounded. Therefore $2\kappa - N$ and T^2/N are integrable and κ/N has a

limit which must be $1/2$ since $\kappa + N \leq 3/\tau + O(\tau)$ due to Prop. 1. This then implies $\tau N \rightarrow 2$ and $\ln r / \ln |\tau| \rightarrow 2$ for $\tau \rightarrow 0$.

With

$$C(\tau) = \int_{\tau}^0 \left(\kappa - \frac{N}{2} \right) d\tau' = O(1), \quad D = e^C r^{\frac{3}{2}}, \quad (74)$$

such that $\dot{D} = (2N - \kappa)D \leq 0$, we then derive

$$W(\tau) = 1 - \int_{\tau}^0 rU d\tau', \quad (75a)$$

$$(rU)(\tau) = \frac{D(\tau)}{D(\tau_0)}(rU)(\tau_0) + \sigma \int_{\tau_0}^{\tau} \frac{D(\tau')}{D(\tau')} W(W^2 - 1) d\tau'. \quad (75b)$$

Starting with $W - 1 = O(1)$, these equations first imply $rU = O(\tau)$ and thus $W - 1 = O(\tau^2)$, and finally $rU = O(\tau^3)$ and thus $W - 1 = O(\tau^4)$. As a consequence U and T are bounded, $\kappa = 1/\tau + O(\tau)$, $N = 2/\tau + O(\tau)$, and $r = -\sigma M \tau^2/2 + O(\tau^3)$ with some $\sigma M < 0$ as required by Eqs. (28,29).

- ii) For κ bounded from below we proceed like in [3], with some modifications since we cannot assume $\kappa \geq 1$. Let A be a positive constant such that $\kappa + A \geq 1$. We introduce $B = e^{A\tau}$ and the new dependent variable $\lambda = W(W^2 - 1) + NU$ and rewrite Eqs. (8c,e) as

$$(BrN)' = (\kappa + A)BrN - 2BrU^2, \quad (76a)$$

$$\dot{U} = \lambda - \kappa U, \quad (76b)$$

$$\dot{\lambda} = (3W^2 - 2U^2 - 1)U. \quad (76c)$$

We want to show, that the r.h.s. of these equations stay bounded as $\tau \rightarrow 0$. In the following ϵ is an arbitrary, suitably small positive number. Since $(BrN)' < 0$ we see that r^ϵ and $|BrN|^{-\epsilon}$ are bounded and monotonic and therefore their derivatives are absolutely integrable at $\tau = 0$. In particular we find

$$\int_0^0 r^\epsilon |N| d\tau < \infty, \quad \int_0^0 \frac{1}{r^{1-\epsilon}} d\tau < \infty, \quad \int_0^0 \frac{U^2}{(Br)^\epsilon |N|^{1+\epsilon}} d\tau < \infty. \quad (77)$$

Next we want to show that $W|BrN|^{-\epsilon}$ is bounded. In view of Prop. 4 this is obvious except when $W^2 \geq 1$ and $W\dot{W} \geq 0$. In that case we get, using the Schwarz inequality, $\int_0^0 |U||BN|^{-\epsilon} d\tau < \infty$ and hence $\int_0^0 |\dot{W}||BrN|^{-\epsilon} d\tau$ is finite for $0 < \epsilon < 1$. Integration by parts yields

$$\begin{aligned} \frac{W}{|BrN|^\epsilon}(\tau) + \epsilon \int_{\tau_0}^{\tau} \frac{W}{|BrN|^\epsilon} \left(\kappa + A - \frac{2U^2}{N} \right) d\tau' \\ = \frac{W}{|BrN|^\epsilon}(\tau_0) + \int_{\tau_0}^{\tau} \frac{\dot{W}}{|BrN|^\epsilon} d\tau', \end{aligned} \quad (78)$$

with some suitably small negative τ_0 . Since the r.h.s. is bounded and both terms on the l.h.s. have the same sign $W|BrN|^{-\epsilon}$ is bounded.

The constraint Eq. (9) implies that $(r(N + A))^2 - (W^2 - 1)^2$ is bounded from above and hence W and with it rN are bounded. As a consequence we get $r(0) = 0$.

Next we use the equation

$$(r^\epsilon U)' = \sigma \frac{W(W^2 - 1)}{r^{1-\epsilon}} + \left((1 + \epsilon)N - \kappa\right)r^\epsilon U. \quad (79)$$

From Lemma 13 we conclude that $r^\epsilon U$ is bounded and consequently $|U|^n$ is integrable for any $n > 0$ on the interval $(\tau_0, 0)$. Hence λ and thus U have a finite limit at $\tau = 0$, implying that also κ is bounded from above. We thus get the RN singularity of Eqs. (22,23). ■

Since Props. 3, 5, and 6 exhaust all possible cases, we obtain the following:

Theorem 7 (Classification for $\sigma = +1$): *A solution of Eqs. (8,9) for $\sigma = +1$ starting at some regular point τ_0 (which is not a f.p.) can be extended in both directions to one of the following singular points:*

- i) a regular origin $r = 0$ at infinite τ as in Eqs. (21);
- ii) a regular horizon of the black hole or cosmological type at some finite τ_1 as in Eqs. (35,36);
- iii) the RN f.p. at infinite τ as in Eqs. (40), where $\kappa_s \leq 0$ for $\tau \rightarrow \pm\infty$ requires $W \equiv 0$;
- iv) $r = 0$ at some finite τ_1 with the S type singularity as in Eqs. (28,29);
- v) $r = 0$ at some finite τ_1 with the RN type singularity as in Eqs. (22,23).

4.2 Solutions with $\sigma = -1$

Due to the presence of the cosmological constant the asymptotics for large r is deSitter instead of Minkowski and this requires $\sigma = -1$. We shall again give a complete classification of all possible solutions starting at some regular point.

Proposition 8: *Suppose a solution of Eqs. (8,9) for $\sigma = -1$ with $N(\tau) \geq 0$ and unbounded, then $r \rightarrow \infty$ for some finite τ_1 with the deSitter asymptotics of Eqs. (37,38).*

Proof: The constraint Eq. (9) implies $2\kappa N \geq N^2 - 1$ and therefore

$$|N| > 1 \quad \text{implies} \quad \kappa N > 0 \quad \text{and} \quad |\kappa| \geq \frac{N^2 - 1}{2|N|} > \frac{|N| - 1}{2}, \quad (80)$$

i.e. κ is unbounded from above. Eq. (56) implies that $(\kappa + N)/r$ is bounded from above and therefore $r \rightarrow \infty$. We then introduce $\tilde{N} = N/r$ and $\tilde{\kappa} = \kappa/r$ and see from the constraint Eq. (9) that $\tilde{N}$ and $\tilde{\kappa}$ are bounded from below by positive constants. This implies that $\tau \rightarrow \tau_1$ stays finite as $r \rightarrow \infty$. In order to be able to use E/r^4 as a Lyapunov function, we introduce a new independent

variable ρ with $d\rho = r d\tau$ such that $\rho \rightarrow \infty$ for $r \rightarrow \infty$, denote the derivative with respect to ρ by a prime, and obtain

$$\left(\frac{E}{r^4}\right)' = -\frac{\bar{\kappa}U^2 + \bar{N}T^2}{r^2}. \quad (81)$$

According to Lemma 11.1 of [19] $(E/r^4)'$ vanishes on the limit points of the solution, showing that U/r and W/r tend to zero for $\rho \rightarrow \infty$. For $\bar{N}$ we find

$$\bar{N}' = \frac{1}{2} \left(\Lambda - 3\bar{N}^2 - \frac{1}{r^2} + \frac{T^2 - 2U^2}{r^2} \right). \quad (82)$$

Applying Lemma 14 we obtain $\bar{N} \rightarrow \sqrt{\Lambda/3}$ and further from Eq. (9) $\bar{\kappa} \rightarrow 2\sqrt{\Lambda/3}$.

Using $s = 1/r$ as independent variable and introducing the new dependent variables $y_\epsilon = (W_\epsilon, \bar{U}_\epsilon, m_\epsilon)$ where $W_\epsilon = s^\epsilon W$, $\bar{U}_\epsilon = s^{2\epsilon} UN$, $m_\epsilon = s^{3\epsilon} m$, and $0 \leq \epsilon \leq 1$, we can rewrite Eqs. (38) as

$$s \frac{d}{ds} W_\epsilon = \epsilon W_\epsilon - s^{1-\epsilon} \frac{\bar{U}_\epsilon}{\bar{N}^2}, \quad (83a)$$

$$s \frac{d}{ds} \bar{U}_\epsilon = 2\epsilon \bar{U}_\epsilon + s^{1-\epsilon} \left(W_\epsilon (W_\epsilon^2 - s^{2\epsilon}) + s^{3(1-\epsilon)} \frac{\bar{U}_\epsilon^3}{\bar{N}^4} \right), \quad (83b)$$

$$s \frac{d}{ds} m_\epsilon = 3\epsilon m_\epsilon + s^{1-\epsilon} \left(\frac{\bar{U}_\epsilon^2}{\bar{N}^2} - \frac{(W_\epsilon^2 - s^{2\epsilon})^2}{2} \right), \quad (83c)$$

with $\bar{N}^2 = \Lambda/3 - s^2 + 2s^{3(1-\epsilon)}m_\epsilon$. The dynamical system Eqs. (83) has three stable modes with real eigenvalues $\geq \epsilon$. Thus $s^{-\eta}y_\epsilon$ is bounded near $s = 0$ for any solution vanishing at $s = 0$ and $\eta < \epsilon$ (see e.g. [18]). Assuming that y_ϵ is bounded and vanishes at $s = 0$ for some $\epsilon = \epsilon_0 > 0$, which we know to be true for $\epsilon_0 = 1$, we therefore conclude that the same is true for all $\epsilon > 2\epsilon_0/3$, and thus by induction for all $\epsilon > 0$. For $\epsilon < 1/4$ we can finally convert Eqs. (38) into integral equations with convergent integrals, i.e. W , UN , and m are bounded as required by Eqs. (37). ■

Proposition 9: Suppose a solution of Eqs. (8,9) for $\sigma = -1$ with N and κ bounded for all $\tau \geq \tau_0$, then the solution runs into

- i) the SdS f.p. described by Eqs. (39) (with $\kappa_s = -1$ only possible if $W^2 \equiv 1$);
- ii) the RN f.p. described by Eqs. (40).

Proof: The constraint Eq. (9) implies that r , U , and T and thus W are bounded. Moreover r has a limit $r_1 \geq 1/\sqrt{2\Lambda}$ since otherwise $\kappa + N$ would diverge due to Eq. (56). Prop. 2 then implies that the solutions runs into

- i) the SdS f.p., as discussed in Section 3. For reasons given there the case $\kappa_s = -1$ is only possible if $W^2 \equiv 1$.
- ii) the RN f.p., as discussed in Section 3. ■

Proposition 10: Suppose a solution of Eqs. (8,9) for $\sigma = -1$ with N unbounded and $N(\bar{\tau}_0) < 0$ for some $\bar{\tau}_0$, then the solution has an origin ($r = 0$) at some finite τ_1 . Depending on the behaviour of N/κ as $\tau \rightarrow \tau_1$ the solution has

- i) an S singularity as described in Eqs. (29), if $N/\kappa \rightarrow 2$;
- ii) a pseudo-RN singularity as described in Eqs. (25), if $N/\kappa \rightarrow 1/2$;
- iii) unbounded oscillating N , U , and T as described in [16], if N/κ has no limit.

Proof: From Eq. (80) we find that $\kappa + N$ is unbounded from below and therefore diverges to $-\infty$ at some finite τ_1 due to Eq. (56). Without restriction we may assume $\tau_1 = 0$. Moreover Eq. (80) implies

$$\kappa < \frac{\kappa + N + 1}{3} \quad \text{for } \kappa + N < -1, \quad (84)$$

and consequently κ diverges as well. Lemma 14 applied to Eq. (8d) then yields $\kappa \leq 1/\tau + O(\tau)$. In the following we can assume $\kappa < -1$, restricting $|\tau|$ to sufficiently small values.

Next we introduce new dependent variables $n = N/\kappa$, $u = U/\kappa$, and $t = T/\kappa$ and note that n , u , and t are bounded due to Eqs. (84) and (9). If n has a limit $n_1 > 0$ then $\int_{\tau_0}^0 N d\tau = -\infty$ and $r \rightarrow 0$.

- i) First consider the case that $n \rightarrow 2$ as $\tau \rightarrow 0$, and consequently $u \rightarrow 0$ and $t \rightarrow 0$ due to the constraint Eq. (9). Prop. 1 then implies $\tau\kappa \rightarrow 1$, $\tau N \rightarrow 2$, and $\ln r / \ln |\tau| \rightarrow 2$ as $\tau \rightarrow 0$. This implies $r\kappa \rightarrow 0$, $rU \rightarrow 0$, and $W^2 \rightarrow 1$ and we may assume $W \rightarrow +1$ without restriction.

Defining

$$C(\tau) = \int_{\tau_0}^{\tau} \left(\frac{N}{2} - \kappa \right) d\tau', \quad D = e^C r^{\frac{3}{2}}, \quad (85)$$

such that $r^\epsilon e^C$ is bounded for $\epsilon > 0$ (compare Eq. (74)), we proceed as in the proof of Prop. 6 and find that the solution has all properties required by Eqs. (28,29).

- ii) Next consider the case that n has a limit $n_1 < 2$. We introduce a new independent variable ρ with $d\rho = |\kappa|d\tau$ such that $\rho \rightarrow \infty$ for $\tau \rightarrow 0$ and denote the derivative with respect to ρ by a prime. We then obtain again Eqs. (30) with $z = 1/|\kappa|$.

First assume $n_1 = 0$ and therefore $u \rightarrow 0$ and $t \rightarrow 0$. Eqs. (30c,e) then first imply that $e^\rho z$ has a finite non-vanishing limit and that $e^\rho t$ and thus T are bounded. Eqs. (30b,d) then imply that $e^{2\rho}u$ and $e^{2\rho}n$ are bounded, i.e. $N \rightarrow 0$ contrary to the assumption that N is unbounded.

For $0 < n_1 < 2$ we see from the constraint Eq. (9) that $t^2 + 2u^2$ has the limit $n_1(2 - n_1) > 0$. From Eq. (30b) we see that for any $\epsilon > 0$ and ρ sufficiently large u is monotonic as long as $|2u^2 - n_1| > \epsilon$ and $|u| > \epsilon$. Since u is bounded it therefore has a limit u_1 with $(2u_1^2 - n_1)u_1 = 0$

and thus t has a limit t_1 . From Eq. (30d) we then obtain the condition $2u_1^2(n_1 + 1) = n_1(2 - n_1)$. This only leaves the possibility $n_1 = |u_1| = |t_1| = 1/2$. Prop. 1 then implies $\tau\kappa \rightarrow 2$, $\tau N \rightarrow 1$, and $\ln r / \ln |\tau| \rightarrow 1$ as $\tau \rightarrow 0$. Thus $r|U|$ is integrable and W has a limit W_1 , and consequently $-rN$, $-r\kappa/2$, and $|rU|$ all have the limit $|W_1^2 - 1|$.

Integrating Eq. (8b) we then find $W = W_1 + O(\tau)$ and therefore $W_1^2 \neq 1$ since otherwise $t \rightarrow 0$ for $\tau \rightarrow 0$. Thus the solution has a pseudo-RN singularity as described in Eqs. (25).

- iii) Finally assume that n has no limit. Since n is bounded this implies that n has an infinite sequence of maxima and minima where the r.h.s. of Eq. (30d) changes sign. Therefore n , u , and t vary between their respective maxima and minima.

If $\int_\rho^\infty n d\rho'$ is finite then the r.h.s. of Eq. (30d) is integrable since $u^2 < n(2 - n) + z^2$ due to the constraint Eq. (9), and therefore n has a limit contrary to the assumption. Thus $\int_\tau^0 N d\tau' = -\int_\rho^\infty n d\rho'$ diverges and $r \rightarrow 0$. ■

Remark: The proof of Prop. 10 also applies to the case $\Lambda = 0$ corresponding to Bartnik-McKinnon type black holes inside their horizon discussed in [16]. In fact, this is to be expected, because the cosmological constant becomes less relevant as r decreases.

Since Props. 3 and 8–10 exhaust all possible cases, we obtain the following:

Theorem 11 (Classification for $\sigma = -1$): *A solution of Eqs. (8,9) for $\sigma = -1$ starting at some regular point τ_0 (which is not a f.p.) can be extended in both directions to one of the following singular points:*

- i) $r = \infty$ at some finite τ_1 with the deSitter asymptotics of Eqs. (37,38);
- ii) a regular horizon of the black hole or cosmological type at some finite τ_1 as in Eqs. (35,36);
- iii) the SdS f.p. at infinite τ as in Eqs. (39), where $\kappa_s = \mp 1$ for $\tau \rightarrow \pm\infty$ requires $W^2 \equiv 1$;
- iv) the RN f.p. at infinite τ as in Eqs. (40);
- v) $r = 0$ at some finite τ_1 with the pseudo-RN type singular behaviour of Eqs. (25,27);
- vi) $r = 0$ at some finite τ_1 with the S type singular behaviour of Eqs. (28,29);
- vii) $r = 0$ at some finite τ_1 with unbounded oscillating N , U , and T as described in [16].

The cases ii), v), and vi) correspond to the NARN, NAPRN, and NAS solutions of [16].

5 4d Space-Times

In the previous section we have determined all possible 1d radial solutions extending between singular points. Taking into account the time independence of the solutions this gives us pieces of 2d space-times with a metric as given in Eq. (2). Our aim in this section is to determine all inextensible (in the sense of geodesic completeness) 4d space-times, which can be constructed from these 2d pieces. A standard way for that is to construct the Carter-Penrose diagrams of these extensions. Recipes for their construction are given in [23, 24]. As a first step, we have to bring the 2-metric to the required standard form

$$ds_2^2 = F dt^2 - \frac{1}{F} d\hat{r}^2, \quad (86)$$

by the coordinate change $d\hat{r} = r e^\nu d\tau$, where $F = \sigma e^{2\nu}$ in terms of our variables. Introducing Eddington-Finkelstein type coordinates via $du_\pm = dt \pm F^{-1} d\hat{r}$ this takes the form

$$ds_2^2 = F du_+ du_-, \quad (87)$$

useful for a Kruskal type extension.

The next step is to express F as a function of $\hat{r}$ and study its behavior near the singular points. This will be done one by one:

- At $r = 0$ the solutions are geodesically incomplete. The singular solutions have a genuine curvature singularity and do not allow for an extension. Although solutions with a regular origin are easily extended taking into account the angular dependence, we treat them as inextensible in the 2d setting.
- At a regular horizon the space-time is geodesically incomplete in the space-like as well as in the light-like direction and F has a simple zero at $\hat{r} = \hat{r}_h$.
- At the RN and SdS fixed points the space-time is always inextensible (i.e. geodesically complete) in the space-like direction. While for $\kappa_s \geq 0$ the coordinate $\hat{r}$ tends to infinity for $\tau \rightarrow \pm\infty$ it stays finite for $\kappa_s \leq 0$. Since $\hat{r}$ plays the role of an affine parameter for light-like geodesics, the space-time is geodesically complete in the first case and incomplete in the second one. For $\kappa\tau \rightarrow -\infty$ one easily finds $\hat{r} \approx \hat{r}_s + e^{\kappa_s \tau} r_s / \kappa_s$ and since $e^{2\nu} \approx e^{2\kappa_s \tau}$ we get

$$F = \sigma e^{2\nu} \approx \sigma \frac{\kappa_s^2}{r_s^2} (\hat{r}_s - \hat{r})^2 \quad (88)$$

near the f.p. This shows that for $\kappa_s \leq 0$ the space-time has a degenerate horizon.

There is an exceptional situation of the RNdS f.p. for $\kappa_s = 0$, which is only possible for $\Lambda = 1/4$, where F has a cubic zero at $r = \sqrt{2}$. Again one has to distinguish the cases $\kappa \geq 0$ and $\kappa \leq 0$ for $\tau \rightarrow \pm\infty$, leading to geodesic completeness resp. incompleteness at this singular point. For $\sigma = +1$ only the RNdS solution with $W \equiv 0$ is possible, but for $\sigma = -1$ there are solutions with non-trivial W . They are C^∞ , but have an essential singularity at the degenerate horizon.

- For solutions with dS asymptotics the space-time is geodesically complete at $r = \infty$.

Before constructing the Carter-Penrose diagrams we present in Figs. 6 and 7 a graphical representation of all possible compositions of the pieces of 2d space-times discussed in the previous section. In principle there is an infinite number of such possible compositions, because one can always insert additional pairs of regular horizons. We have, however, restricted ourselves to compositions which are compatible with the number of parameters of a generic solution. Thus we take into account the necessity to suppress the divergent modes at every singular point. For example at every horizon we loose one parameter. In this counting the singular solutions at $r = 0$ are considered as f.p.s without divergent mode (compare Sect. 3) as are the solutions with dS asymptotics. For a given Λ the five autonomous Eqs. (8) are reduced to four taking into account the constraint Eq. (9). This results in a 3-parameter family of orbits. Hence we can suppress maximally four divergent modes varying also Λ , implying e.g. that we can have a maximum of four horizons. This does, however, not necessarily imply that all cases are realized, i.e. can be actually obtained numerically. Nor does it mean that solutions with more than four horizons are excluded. It just means that we see no way to construct them varying the available parameters at hand. Actually we have not even found any solution with four horizons.

The graphs in Figs. 6 and 7 are more or less self explanatory, but some remarks are in order. We ignore the presence resp. position of an equator, which would just lead to a much larger number of diagrams without a particular gain in information. Likewise we ignore the type of the singularity at a singular origin.

Diagram 2 represents the generic SdS solution, while 3 corresponds to the degenerate case. Diagram 5 is the case of SdS with negative mass. Diagram 6 is the generic RNdS solution, 9 corresponds to the most degenerate RNdS solution for $\Lambda = 1/4$ with a cubically degenerate horizon. Diagrams 4, 10, and 16 represent generic solutions. The space-times 10–23 without horizons either have spatial sections with 3-sphere topology or have infinite throats with dS resp. AdS type asymptotic behaviour. Diagrams 3 and 8 represent the degenerate SdS resp. RNdS solutions as well as ones with non-trivial YM field. Diagram 9 describes besides the most degenerate RNdS solution also solutions joining the $\sigma = +1$ piece of the latter to a non-trivial $\sigma = -1$ part at the cubically degenerate horizon. The same property is shared by the diagrams 32, 33, and 42.

Diagrams 15, 22, and 23 describe the degenerate space-time of Eqs. (20a). Diagram 41 describes a solution with a degenerate horizon running into the SdS f.p. Such a solution could occur as a limit of a 1-parameter family of solutions of the type 8 with YM hair.

The solutions with a regular origin of [9] are given by diagrams 1, 28, 29, and 13 in the order as they occur with increasing Λ . Replacing the regular origin by a regular black hole horizon one obtains the diagrams 2, 36, 34, and 29. Replacing the origin by two horizons as in the RN black hole one obtains the diagrams 6, 44, 43, and 39. The diagrams 19, 20, and 21 are new types of regular solutions starting at the RN resp. SdS f.p. and running into

Table 1: Parameters of regular solutions

n	Λ_c b_c	r_c W_c	Λ_* b_*	r_* W_*	Λ_r b_r
1	0.33049695	1.729083	0.33448289	1.706678	0.75000000
	0.37534807	-0.85135	0.37417940	-0.85218	0.25000000
2	0.23896851	1.934703	0.25004507	1.812673	0.36424423
	0.54743193	+0.45963	0.53831284	+0.48411	0.42959769
3	0.23712384	1.832376	0.24706870	1.686850	0.29321764
	0.57517232	-0.23626	0.56568105	-0.28415	0.50882906
4	0.24027732	1.746936	0.24790151	1.606049	0.27032753
	0.57699432	+0.13396	0.56950069	+0.18872	0.54048970
5	0.24275653	1.688873	0.24861452	1.556666	0.26089512
	0.57578215	-0.08158	0.56992909	-0.13598	0.55401931
6	0.24445557	1.648272	0.24907014	1.524611	0.25638498
	0.57448821	+0.05216	0.56981254	+0.10345	0.56042575

the SdS f.p. or to infinity.

Figs. 8–13 show the elementary building blocks of the Carter-Penrose diagrams corresponding to the diagrams of Figs. 6 and 7, which have at least one horizon. The complete diagrams are obtained from those by periodic repetition, gluing them along suitable common horizons [23, 24]. As usual space-like boundaries are represented by horizontal lines, time-like ones are vertical.

6 Numerical Results

In this section we present a selection of numerical results, some of which we would like to emphasize. We perform a quantitative study of the limit $n \rightarrow \infty$ of the 3-sphere solutions. We present a whole new class of solutions with two and three regular horizons. We give a complete description of the phase space of solutions for $\sigma = +1$ with a regular origin. An analogous description for $\sigma = -1$ exhibits some important aspects. One cannot expect a complete description due to the very complicated structure of solutions with a singular origin caused by unbounded oscillations [15, 16].

6.1 Solutions with a regular origin and a regular horizon

In Fig. 14 we show parameters of solutions with a regular origin and a regular horizon with $n = 1$ and 2 zeros of W as well as for the limiting solutions ($n = \infty$) connecting the origin and the RN f.p. In Table 1 we present parameters of these solutions for $n \leq 6$ and some special values of Λ . When continued beyond the horizon into the region with $\sigma = -1$ they are asymptotically deSitter for Λ up to an n -dependent critical value Λ_c . For $\Lambda = \Lambda_c$, indicated by the points C_n in Fig. 14 they run into the SdS f.p. while for $\Lambda > \Lambda_c$ they have an equator and

Table 2: Parameters of 3-sphere solutions

n	Λ	b	$r_{\max}$
1	0.7500000000	0.2500000000	1.4142136
2	0.3642442330	0.4295976874	1.4355566
3	0.2932176375	0.5088290567	1.5213897
4	0.2703275292	0.5404896987	1.4826529
5	0.2608951239	0.5540193079	1.4829530
10	0.2512790802	0.5674090979	1.4396422
20	0.2501165095	0.5688910784	1.4224344
30	0.2500264064	0.5690006373	1.4181848
40	0.2500089730	0.5690215671	1.4165412
50	0.2500038394	0.5690276981	1.4157401
∞	0.25	0.5690322659	$\sqrt{2}$

end at $r = 0$. The equator coincides with the horizon for $\Lambda = \Lambda_*$ (the open circles in Fig. 14) and occurs for $\sigma = \mp 1$ if $\Lambda \leq \Lambda_*$. The radius of the horizon decreases as Λ increases and vanishes for $\Lambda = \Lambda_r$, the points R_n , corresponding to the regular 3-sphere solutions.

6.2 Regular 3-sphere solutions

Table 2 contains the parameters of some regular 3-sphere solutions with up to 50 zeros of W . As n increases the values Λ_n , b_n rapidly converge to $\Lambda_\infty = 1/4$ and $b_\infty \approx 0.569032$, i.e. the point R_∞ of Fig. 14. The corresponding limiting solution connects a regular origin and the RN f.p. with $\Lambda = 1/4$ and asymptotic behaviour (compare Sect. 3.8).

$$\kappa \approx \frac{4}{\tau - \tau_0}, \quad W \approx \sqrt{84}r \frac{\sin(\tau + \theta)}{(\tau - \tau_0)^2}. \quad (89)$$

Numerically we find $\theta \approx 1.7884$ and $\tau_0 \approx -3.8717$ (normalizing τ such that $\tau - \ln r \rightarrow 0$ as $r \rightarrow 0$). Solutions with $(\Lambda_\infty + \delta\Lambda, b_\infty + \delta b)$ for small $\delta\Lambda$, δb will miss the RN f.p. but yield a regular 3-sphere provided κ , N , and W or U simultaneously vanish for some τ . Assuming that the phase of W remains essentially unchanged this requires that κ and N vanish at $\tau + \theta \approx (n+1)\pi/2$ in order to have n zeros of W . For solutions starting from the origin we first have to choose $\delta b \approx -1.2\delta\Lambda$ to suppress the unstable mode (with eigenvalue +6) near the f.p., where the solution approximately obeys

$$\dot{\kappa} = \bar{r} - \kappa^2, \quad (90a)$$

$$\dot{\bar{r}} = -2\bar{N}, \quad (90b)$$

$$\dot{\bar{N}} = -\frac{\bar{r}^2}{4} - \delta\Lambda, \quad (90c)$$

with the consequence that $\bar{r}^3/12 + \delta\Lambda\bar{r} - \bar{N}^2$ is constant. Moreover Eqs. (90) are invariant under the rescaling $(d\tau, \kappa, \bar{r}, \bar{N}, \delta\Lambda) \rightarrow (d\tau/\lambda, \lambda\kappa, \lambda^2\bar{r}, \lambda^3\bar{N}, \lambda^4\delta\Lambda)$.

Starting from some τ_1 with $\kappa = \bar{N} = 0$ and $\bar{r} = -1$ we can thus integrate

$$\dot{\kappa} = \bar{r} - \kappa^2, \quad (91a)$$

$$\dot{\bar{r}} = -\sqrt{\frac{\bar{r}^3 + 1}{3} + 4\delta\Lambda(\bar{r} + 1)}, \quad (91b)$$

towards decreasing τ , adjusting τ_1 and $\delta\Lambda$ such that κ and $\bar{r}$ diverge at $\tau = \tau_0$ with $\bar{r}/\kappa^2 \rightarrow 3/4$. Numerically we find $\delta\Lambda \approx 0.81575$ and $\tau_1 - \tau_0 \approx 3.8288$ and thus obtain for Λ_n and $\bar{r}_n$, the value of $\bar{r}$ at the equator, the asymptotic expressions

$$\left((n+1)\frac{\pi}{2} - \theta - \tau_0\right) \sqrt[4]{\Lambda_n - 0.25} \approx 3.8288 \sqrt[4]{0.81575} \approx 3.6387, \quad (92a)$$

$$\frac{\bar{r}_n}{\sqrt{\Lambda_n - 0.25}} \approx -\frac{1}{\sqrt{0.81575}} \approx -1.1072, \quad (92b)$$

in good agreement with the numerical results shown in Fig. 15 for $n \leq 50$.

6.3 Solutions with two or more regular horizons

Fig. 16 shows parameters for solutions with two horizons for $\sigma = +1$ and various values of Λ . In the limit $r_h \rightarrow 0$ for one or both horizons they reproduce the solutions with an origin and a horizon and the regular 3-spheres discussed above. The dashed-dotted curve delimits the region of existence; on it the value of W_h vanishes and the solutions bifurcate with Reissner-Nordström-deSitter solutions. The dotted curves correspond to the critical Λ values, giving solutions running into the Schwarzschild f.p. in the region $\sigma = -1$. The solutions in the region outside the dotted curves are asymptotically deSitter. The dashed curves correspond to solutions with coinciding equator and horizon. Solutions in the region inside the dashed curves have an equator between their horizons in contrast to those outside of it. The RN-deSitter and the critical solutions have no equator and thus the dashed-dotted and the dotted curves must lie outside the dashed ones. Since on the diagonal the two horizons are equal, these solutions have an equator and thus the diagonal lies inside the dashed curves. Thus the dashed curves must go through the point P on the intersection of the diagonal with the boundary curve, where the two horizons merge. Since the solutions on the boundary curve are asymptotically deSitter, the dotted curve must also go through P . At this point the solution can be obtained exactly through a suitable rescaling. Putting

$$W = \epsilon \bar{W}, U = \epsilon \bar{U}, N = \epsilon^2 \bar{N} \quad \text{and} \quad r = R(1 + \epsilon^2 \bar{r}), \quad (93)$$

where ϵ is a small parameter we obtain in leading order from the constraint Eq. (9) $\Lambda = (R^2 - 1)/r^4$ and from Eqs. (8)

$$\dot{\bar{W}} = R\bar{U} \quad (94a)$$

$$\dot{\bar{U}} = -\frac{\bar{W}}{R} - \kappa \bar{U} \quad (94b)$$

$$\dot{\kappa} = 1 - 2\Lambda R^2 - \kappa^2. \quad (94c)$$

Table 3: Some solutions with three regular horizons and two zeros of W

Λ	W_0	r_1	W_1	r_2	W_2	r_3	W_3	W_∞
0.0045	1.3655	0.0948	1.3582	1.0256	0.1000	24.6488	0.0862	0.1637
0.0643	1.4072	0.1207	1.3964	1.0459	0.0800	5.6355	0.0263	0.0638
0.1187	1.4388	0.1417	1.4249	1.0918	0.0600	3.7151	0.0197	0.0610
0.1655	1.4737	0.1664	1.4556	1.1502	0.0400	2.8234	0.0152	0.0647
0.2061	1.5141	0.1968	1.4904	1.2351	0.0200	2.2350	0.0098	0.0716
0.2247	1.5378	0.2155	1.5103	1.3051	0.0100	1.9718	0.0059	0.0762
0.2337	1.5511	0.2263	1.5214	1.3623	0.0050	1.8255	0.0034	0.0789
0.2411	1.5630	0.2361	1.5312	1.4544	0.0010	1.6627	0.0008	0.0813

Solving Eq. (94c) we get $\kappa = a \cot(a\tau)$ as in Eq. (20c) with $a^2 = 2\Lambda R^2 - 1$ and plugging the result into Eqs. (94a,b) we arrive at the second order equation

$$\ddot{\bar{W}} + a \cot(a\tau) \dot{\bar{W}} + \bar{W} = 0 . \quad (95)$$

Introducing $x = \cos(a\tau)$ we find the Legendre equation

$$\frac{d}{dx} \left((1-x^2) \frac{d\bar{W}}{dx} \right) + \frac{1}{a^2} \bar{W} = 0 . \quad (96)$$

The acceptable solutions are those regular at $x = \pm 1$ given by the Legendre polynomials, obtained for $a^{-2} = n(n+1) = 2, 6, 12, \dots$ yielding

$$R^2 = \frac{2}{1-a^2} = 4, \frac{12}{5}, \frac{24}{11}, \dots , \quad (97a)$$

$$\Lambda = \frac{1-a^4}{4} = \frac{3}{16}, \frac{35}{144}, \frac{143}{576}, \dots . \quad (97b)$$

The cases $n = 1$ and 2 are shown in Fig. 16.

Combining numerical results for $\sigma = +1$ and $\sigma = -1$ we have found a one parameter family of solutions with three regular horizons and two zeros of W . In Table 3 we present the parameters of some of these solutions. Starting from $r = 0$ with an RN type singularity and $W = W_0$, they have two regular horizons at $r = r_{1,2}$ with $W = W_{1,2}$, two zeros of W , a third horizon at r_3 with W_3 , and extend to $r = \infty$ with dS asymptotics and $W = W_\infty$. There is a similar such family with three zeros of W , but we found no such solution with only one zero.

6.4 Description of the phase space

Fig. 17 schematically shows the phase space for solutions starting from a regular origin, i.e. with $\sigma = +1$, dividing the (Λ, b) plane into various regions according to their type and including results for $\Lambda = 0$ from [3] resp. for $\Lambda < 0$ from [12]. All the cases listed in the Classification Thm. 7 are present in the half plane $\Lambda > 0$ of Fig. 17. The regions AdS_n with $\Lambda < 0$ and S_n correspond to solutions with n zeros of W and AdS asymptotics or a singular origin respectively. At the moment this picture is mostly based on numerical results, but proving the

existence of such a picture will certainly be an essential part in an existence proof for the various solution types.

In order to simplify the description of the phase space we deform and compactify Fig. 17 into Fig. 18 as follows: First we identify the points $(\Lambda, \pm\infty)$ and add a dashed-dotted line along $b = \pm\infty$ separating the two previously disconnected parts of the region S_0 . Next we add a point $R_0 = (+\infty, 0)$ corresponding to the limit $r_h \rightarrow 0$ of deSitter space and compactify $(+\infty, b)$ to this point. The points with constant (finite) Λ are then on concentric curves around R_0 . We then add the points b_{-1} , P_{-1} , and R_{-1} as the limits $\Lambda \rightarrow -\infty$ of $(\Lambda, -\sqrt{|\Lambda|/4})$, $(\Lambda, +\sqrt{|\Lambda|/4})$, and $(\Lambda, \pm\infty)$. And we add points P_n for $n = 1, \dots$ all identical to P_∞ . Finally the boundary of Fig. 18 corresponding to $(-\infty, b)$ is compactified to one point $b_{-1} = P_{-1} = R_{-1}$.

The open region $\text{AdS} = \cup_{n \geq 0} \text{AdS}_n$ is the complement of the closure of the open region $S = \cup_{n \geq 0} S_n$. The points b_n at $(0, b_n)$ represent the asymptotically flat BK solutions (augmented by Minkowski space for $n = 0$), with their limit point b_∞ (compare [3]). The points R_n represent the regular 3-sphere solutions, with their limit point R_∞ at $\Lambda = 1/4$. AdS_n and AdS_{n+1} are separated by the dotted curves from b_n to P_n where $W_\infty = 0$; S_n and S_{n+1} are separated by the dotted curves from R_n to P_n where $W_0 = 0$ and the solid curves from b_n to R_n with a regular horizon. The dashed-dotted curves from R_{n-1} to R_n with $|W_0| = 1$ and an S type singularity at the origin divide S_n into parts with $|W_0| > 1$ and $|W_0| < 1$, both with a RN type singularity. Similar dashed-dotted curves from b_{n-1} to b_n where $|W_\infty| = 1$ separate AdS_n into parts with $|W_\infty| > 1$ and $|W_\infty| < 1$. The regions AdS_n and S_n are separated by the solid curves from b_{n-1} to b_n with SAdS asymptotics where W_∞ resp. W_0 diverge (compare [12]). Finally the dashed curve from R_∞ through b_∞ , P_∞ with $\Lambda = -15/4$, and P_0 to P_{-1} represents solutions with the RN f.p. asymptotics. They oscillate for $\Lambda > -15/4$ and therefore all regions AdS_n and S_n for $n \gg 1$ are in the vicinity of the curve from R_∞ to P_∞ . For $\Lambda < -15/4$ the RN solutions have a finite number of zeros; numerically we found $n = 1$ from P_∞ to P_0 with $\Lambda \approx -5.0646$ and $n = 0$ from P_0 to P_{-1} . The curves of constant W_∞ in the region AdS_n all connect b_{n-1} and b_n whereas the curves of constant W_0 in the region S_n all connect R_{n-1} and R_n , approaching the solid boundary as W_∞ resp. W_0 diverges or the dotted and dashed one as W_∞ resp. W_0 vanishes.

Fig. 19 shows the phase space of solutions for $\sigma = -1$ starting from a regular horizon with at $r_h = 0.1$, again with all the cases listed in the Classification Thm. 11. The open region $\text{dS} = \cup_{n \geq 0} \text{dS}_n$ of the (Λ, W_h) plane corresponds to solutions with dS asymptotics; this region is divided into regions dS_n with n zeros of W , separated by the dashed-dotted curves where $W_\infty = 0$. The boundary of dS represents solutions ending in a f.p. with finite r : the SdS f.p. with $W = (-1)^n$ at the solid curves bounding dS_n or the RN f.p. with $W = 0$ at the points Q_i connecting these curves.

The complement of the closure of dS represents different types of solutions. They can end at a second regular horizon (dashed curves) or a singular origin with either an S type singularity (dotted curves), a pseudo-RN type singularity (cross at $(\Lambda, W_h) \approx (0.09, 0.5)$), or unbounded oscillations. In Fig. 19 only a selection of such curves is shown.

7 Appendix

First we extend Prop. 1 of [3] by a uniqueness statement:

Proposition 12: *Suppose a system of differential equations for $m+n$ functions $u = (u_1, \dots, u_m)$ and $v = (v_1, \dots, v_n)$,*

$$t \frac{du_i}{dt} = t^{\mu_i} f_i(t, u, v), \quad t \frac{dv_i}{dt} = -\lambda_i v_i + t^{\nu_i} g_i(t, u, v), \quad (98)$$

with constants λ_i with $\operatorname{Re}(\lambda_i) > 0$ and integers $\mu_i, \nu_i \geq 1$ and let $\mathcal{C}$ be an open subset of R^m such that the functions f and g are analytic in a neighbourhood of $t = 0$, $u = c$, $v = 0$ for all $c \in \mathcal{C}$. Then there exists a unique m -parameter family of solutions of the system (12) with boundary conditions $(u, v) = (c, 0)$ at $t = 0$, defined for $c \in \mathcal{C}$ and $|t| < t_0(c)$ with some $t_0(c) > 0$. These solutions satisfy

$$u_i(t) = c_i + O(t^{\mu_i}), \quad v_i(t) = O(t^{\nu_i}), \quad (99)$$

and are analytic in t and c .

Proof: The proof of existence and analyticity in [3] for real $\lambda_i > 0$ applies without any change to complex values with positive real part.

In order to prove uniqueness we assume (without loss of generality) $t \geq 0$, introduce $\tau = -\ln t$ as new independent variable as well as $s = \sqrt{t}$ as additional dependent variable, and replace Eqs. (98) by the autonomous system

$$\dot{s} = -\frac{s}{2}, \quad (100a)$$

$$\dot{\bar{u}}_i = \frac{\bar{u}_i}{2} - s^{2\mu_i-1} f_i(s^2, c + s\bar{u}, v), \quad (100b)$$

$$\dot{v}_i = \lambda_i v_i - s^{2\nu_i} g_i(s^2, c + s\bar{u}, v), \quad (100c)$$

where $\bar{u}_i = (u_i - c_i)/s$ vanishes as $\tau \rightarrow \infty$, i.e. $s \rightarrow 0$ due to Eqs. (99). This system has the hyperbolic f.p. $s = \bar{u} = v = 0$ with one stable and $m+n$ unstable modes. Thus there exists a unique 1-dimensional stable manifold with $(\bar{u}, v)$ analytic in s . Since Eqs. (100) are invariant under the substitution $(s, \bar{u}, v) \rightarrow (-s, -\bar{u}, v)$ this implies that (u, v) is analytic in t . ■

Next we prove some simple Lemmata used in the proof of the classification theorems.

Lemma 13: *Consider a solution y of the linear differential equation $\dot{y} = a + by$ in some interval $\tau_0 \leq \tau < \tau_1$ with $|a|$ integrable. If*

$$c(\tau', \tau) = \int_{\tau'}^{\tau} b(\tau'') d\tau'', \quad (101)$$

is bounded from above for $\tau_0 \leq \tau' \leq \tau < \tau_1$ then y is bounded; if $c(\tau', \tau)$ has a limit as $\tau \rightarrow \tau_1$ then $y(\tau)$ has a limit; if $c(\tau', \tau_1) = -\infty$ then $y(\tau_1) = 0$.

Proof: All properties are implied by the explicit form

$$y(\tau) = y(\tau_0) e^{c(\tau_0, \tau)} + \int_{\tau_0}^{\tau} a(\tau') e^{c(\tau', \tau)} d\tau'. \quad (102)$$

■

Lemma 14: Consider solutions of the Riccati equation $\dot{y} = a - y^2$.

- i) Suppose $|a|$ is integrable on some interval $\tau_0 \leq \tau \leq \tau_1$, then the solution which is unbounded at τ_1 behaves as $y = 1/(\tau - \tau_1) + O(1)$ as $\tau \rightarrow \tau_1$. If a is bounded then $y = 1/(\tau - \tau_1) + O(\tau - \tau_1)$.
- ii) Suppose a is bounded from below and $y < 0$ for $\tau_0 \leq \tau \leq \tau_1$ and y diverges as $\tau \rightarrow \tau_1$, then $y \leq 1/(\tau - \tau_1) + O(\tau - \tau_1)$.
- iii) Suppose a has a limit a_1 for $\tau \rightarrow \infty$ and y is bounded, then a_1 must be non-negative and $y(\tau) \rightarrow \pm\sqrt{a_1}$ for $\tau \rightarrow \infty$.

Proof:

- i) Without restriction we assume $\tau_1 = 0$, put $y = 1/\tau + \eta$, and get by integration

$$\eta(\tau) = \int_{\tau}^0 \left(\frac{\tau'}{\tau} \right)^2 \left(\eta(\tau')^2 - a(\tau') \right) d\tau'. \quad (103)$$

which we solve by iteration starting with $\eta_0 = 0$. Putting $A = \int_{\tau_0}^0 (\tau'/\tau)^2 |a(\tau')| d\tau'$ and $\|f\| = \sup\{|f(\tau)|, \tau_0 \leq \tau < 0\}$ we can estimate $\|\eta_n\| \leq A(1 - |\tau_0|/3)^{-1} < 1$ choosing $|\tau_0|$ small enough. From this we get $\|\eta_{n+1} - \eta_n\| \leq (2|\tau_0|/3)^n \|\eta_1\|$ and thus convergence for $|\tau_0| < 3/2$. For $|a| \leq B$ and $\tau_0^2 B \leq 1$ we can estimate $\|\eta_n/\tau\| \leq B/2$.

- ii) Let A be a constant such that $a \geq A$. We assume again $\tau_1 = 0$, put $\eta = 1/(\tau y)$, and get by integration

$$\eta(\tau) = 1 + \int_{\tau}^0 \frac{\tau'^2}{\tau} a \eta^2 d\tau' \leq 1 + A \int_{\tau}^0 \frac{\tau'^2}{\tau} \eta^2 d\tau'. \quad (104)$$

For $A \geq 0$ we obtain immediately $\eta \leq 1$ and thus $y \leq 1/\tau$. For $A < 0$ we choose $|\tau_0|$ sufficiently small such that $|A|\tau_0^2 \leq 1/2$ and can estimate $\eta \leq 1 + |A|\tau^2$ and consequently $y \leq 1/\tau - |A|\tau$.

- iii) Obviously a_1 must be non-negative, otherwise y would be unbounded from below. For any $\epsilon > 0$ we can find τ_ϵ such that $|a - a_1| < \epsilon/2$ for $\tau > \tau_\epsilon$. If $y(\tau_\epsilon) < -\sqrt{a_1 + \epsilon}$, then y will be unbounded from below. Now suppose that $y(\tau_\epsilon) > \sqrt{a_1 + \epsilon}$, then we can find $\tau'_\epsilon > \tau_\epsilon$ such that y decreases for $\tau_\epsilon < \tau \leq \tau'_\epsilon$ and $y < \sqrt{a_1 + \epsilon}$ for $\tau > \tau'_\epsilon$. If $a_1 = 0$ this implies $y \rightarrow 0$ for $\tau \rightarrow \infty$. For $\epsilon < a_1 \neq 0$ now suppose that $y^2(\tau_\epsilon) < a_1 - \epsilon$, then we can find $\tau'_\epsilon > \tau_\epsilon$ such that y increases for $\tau_\epsilon < \tau \leq \tau'_\epsilon$ and $y > \sqrt{a_1 - \epsilon}$ for $\tau > \tau'_\epsilon$. This implies $y \rightarrow \sqrt{a_1}$ for $\tau \rightarrow \infty$. On the other hand, if $-\sqrt{a_1 + \epsilon} < y(\tau_\epsilon) \leq -\sqrt{a_1 - \epsilon}$ for any ϵ then $y \rightarrow -\sqrt{a_1}$. ■

Lemma 15: Consider a solution of the Riccati equation $\dot{y} = a + by - y^2$. If a is bounded from above and b is bounded for $\tau \geq \tau_0$, then y is either bounded for all $\tau \geq \tau_0$ or diverges to $-\infty$ for some finite $\tau_1 > \tau_0$.

Proof: Let A, B be positive constants such that $a < A$ and $|b| < B$. We can estimate $\dot{y} < 0$ for $|y| > C = \sqrt{2A} + 2B$ and therefore y is bounded from above. Furthermore y monotonically decreases and $(1/y)' > 1/2$ for $y < -C$, and thus $y \rightarrow -\infty$ for some finite τ_1 . ■

Acknowledgements

We would like to thank H. Giacomini and M.T. Grau for helpful discussions on dynamical systems. We also thank the Theoretical Physics groups of the Max-Planck-Institute for Physics resp. University of Tours for their kind hospitality during several visits, where part of the work was done.

References

- [1] R. Bartnik and J. McKinnon, *Phys. Rev. Lett.* **61** (1988) 141
- [2] J.A. Smoller and A.G. Wasserman, *Commun. Math. Phys.* **151** (1993) 303
- [3] P. Breitenlohner, P. Forgács and D. Maison, *Commun. Math. Phys.* **163** (1994) 141
- [4] H.P. Künzle and A.K.M. Masood ul Alam, *J. Math. Phys.* **31** (1990) 928
- [5] P. Bizón, *Phys. Rev. Lett.* **64** (1990) 2844
- [6] M.S. Volkov and D.V. Gal'tsov, *JETP Lett.* **50** (1989) 346
- [7] J.A. Smoller, A.G. Wasserman and S.T. Yau, *Commun. Math. Phys.* **154** (1993) 377
- [8] P. Breitenlohner and D. Maison, *Commun. Math. Phys.* **171** (1995) 685
- [9] M.S. Volkov, N. Straumann, G.V. Lavrelashvili, M. Heusler and O. Brodbeck: *Phys. Rev. D* **54** (1996) 7243
- [10] E. Winstanley, *Class. Quantum Grav.* **16** (1999) 1963
- [11] J. Bjoraker, Y. Hosotani, *Phys. Rev. Lett.* **84** (2000) 1853
- [12] P. Breitenlohner, D. Maison and G. Lavrelashvili, *Class. Quantum Grav.* **21** (2004) 1–17
- [13] A.N. Linden, *Commun. Math. Phys.* **221** (2001) 525
- [14] A.N. Linden, ‘A Classification of Spherically Symmetric Static Solutions of SU(2) Einstein Yang Mills Equations with Non-negative Cosmological Constant’, gr-qc/0005006
- [15] E.E. Donets, D.V. Gal'tsov, M.Yu. Zotov, *Phys. Rev. D* **56** (1997) 3459

- [16] P. Breitenlohner, G. Lavrelashvili and D. Maison, *Nucl. Phys.* **B 524** (1998) 427
- [17] H. Nariai, *Science Reports of the Tohoku Univ.* **35** (1951) 62
- [18] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*. New York: McGraw-Hill, 1955.
- [19] P. Hartman, *Ordinary Differential Equations*. Boston: Birkhäuser, 1982.
- [20] V.I. Arnold and Yu.S. Il'yashenko, *Ordinary Differential Equations in Dynamical Systems I*, D.V. Anosov and V.I. Arnold eds., Berlin, Heidelberg: Springer, 1988.
- [21] F. Dumortier, *Techniques in the Theory of Local Bifurcations: Blow-Up, Normal Forms, Nilpotent Bifurcations, Singular Perturbations in Bifurcations and Periodic Orbits of Vector Fields*, D. Schlomiuk ed., NATO ASI Series C: Math and Phys. Sciences **408** (1993) 19–73
- [22] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge University Press, 1973.
- [23] M. Walker, *J. Math. Phys.* **11** (1970) 2280
- [24] M.O. Katanaev, T. Kloesch and W. Kummer, *Annals Phys.* **276** (1999) 191
M.O. Katanaev, *Nucl. Phys. Proc. Suppl.* **88** (2000) 233–236




Figure 1: Solutions near a singular origin with $|W_0| \approx 1$




Figure 2: Four trajectories of Eqs. (32) emerging from the focal point $(\xi, \eta, \zeta) = (1/2, 1, 1/2)$ and rapidly approaching the bounding triangle




Figure 3: Allowed domains C_+ for black hole boundary conditions for $\Lambda = 0.05, 0.15, 0.25, 0.4$, and 1




Figure 4: Solutions near a regular horizon with $r_h \ll 1$




Figure 5: RN f.p. with $\Lambda = 1/4$: Tangential flow after the blow-up




Figure 6: 1d graphs




Figure 7: 1d graphs with horizons




Figure 8: Carter-Penrose diagrams: one regular or degenerate horizon




Figure 9: Carter-Penrose diagrams: two regular horizons




Figure 10: Carter-Penrose diagrams: one degenerate and one regular horizon




Figure 11: Carter-Penrose diagrams: three regular horizons




Figure 12: Carter-Penrose diagrams: one degenerate and two regular horizons




Figure 13: Carter-Penrose diagrams: four regular horizons




Figure 14: Regular solutions with $n = 1, 2, \text{ and } \infty$ zeros of W




Figure 15: Asymptotic behaviour of the parameters of regular 3-sphere solutions




Figure 16: r_{h1} vs. r_{h2} for solutions with $n = 1$ and 2 zeros of W




Figure 17: Schematic phase space of solutions for $\sigma = +1$ with a regular origin




Figure 18: Compactified phase space of solutions for $\sigma = +1$ with a regular origin




Figure 19: Phase space of solutions for $\sigma = -1$ starting from a regular horizon with $r_h = 0.1$